

Fractional Iteration of Series and Transseries

G. A. Edgar

February 11, 2010

Abstract

We investigate compositional iteration of fractional order for transseries. For any large positive transseries T of exponentiality 0, there is a family $T^{[s]}$ indexed by real numbers s corresponding to iteration of order s . It is based on Abel's Equation. We also investigate the question of whether there is a family $T^{[s]}$ all sharing a single support set. A subset of the transseries of exponentiality 0 is divided into three classes ("shallow", "moderate" and "deep") with different properties related to fractional iteration.

Introduction

Since at least as long ago as 1860 (A. Cayley [3]) there has been discussion of real iteration groups (compositional iteration of fractional order) for power series. Or at least for formal power series, where we do not worry about convergence of the result. In this paper we adapt this to **transseries**. In many cases it is, in fact, not difficult to do when we ignore questions of convergence.

We will primarily use the ordered differential field $\mathbb{T} = \mathbb{R}[[\mathfrak{G}]] = \mathbb{R}[[x]]$ of (real grid-based) transseries; $\mathbb{T}$ is also known as **the transline**. So $\mathbb{T}$ is the set of all grid-based real formal linear combinations of monomials from $\mathfrak{G}$, while $\mathfrak{G}$ is the set of all e^L for $L \in \mathbb{T}$ purely large. (Because of logarithms, there is no need to write separately two factors as $x^b e^L$.) See "Review" below.

The problem looks like this: Let T be a large positive transseries. Is there a family $T^{[s]}$ of transseries, indexed by reals s , so that: $T^{[0]}(x) = x$, $T^{[1]} = T$, and $T^{[s]} \circ T^{[t]} = T^{[s+t]}$ for all $s, t \in \mathbb{R}$? These would be called **fractional iterates** of T : $U = T^{[1/2]}$ satisfies $U \circ U = T$; or $T^{[-1]}$ is the compositional inverse of T ; etc. We limit this discussion to *large positive* transseries since that is where compositions $S \circ T$ are always defined. In Corollary 4.5 we conclude (in the well-based case) that any large positive T of exponentiality 0 admits such a family of fractional iterates. However, there are grid-based large positive transseries T and reals s for which the fractional iterate $T^{[s]}$ is not grid-based (Example 4.8).

We also investigate the existence of a family $T^{[s]}, s \in \mathbb{R}$, all supported by a single grid (in the grid-based case) or by a single well-ordered set (in the well-based case). We show that such a family exists in certain cases (Theorems 3.7 and 3.8) but not in other cases (Theorem 3.9).

Review

The differential field $\mathbb{T}$ of transseries is completely explained in my recent expository introduction [9]. Other sources for the definitions are: [1], [5], [8], [15]. I will generally follow the notation from [9]. The well-based version of the construction is described in [8] (or [10, Def. 2.1]). In this paper it is intended that all results hold for both versions, unless otherwise noted. (I use labels $\boxed{\text{G}}$ or $\boxed{\text{W}}$ for statements or proofs valid only for the grid-based or well-based version, respectively.) We will write $\mathbb{T}$, $\mathfrak{G}$, and so on in both versions.

Write $\mathcal{P} = \{S \in \mathbb{T} : S \succ 1, S > 0\}$ for the set of large positive transseries. The operation of composition $T \circ S$ is defined for $T \in \mathbb{T}$, $S \in \mathcal{P}$. The set $\mathcal{P}$ is a group under composition ([15, § 5.4.1], [8, Cor. 6.25], [10, Prop. 4.20], [11, Sec. 8]). Both notations $T \circ S$ and $T(S)$ will be used.

We write $\mathfrak{G}$ for the ordered abelian group of transmonomials. We write $\mathfrak{G}_{N,M}$ for the transmonomials with exponential height N and logarithmic depth M . We write $\mathfrak{G}_N$ for the log-free transmonomials with height N . Even in the well-based case, the definition is restricted so that for any $T \in \mathbb{T}$ there exists N, M with $\text{supp } T \subseteq \mathfrak{G}_{N,M}$. The support $\text{supp } T$ is well ordered for the converse of the relation $\prec$ in the well-based case; the support $\text{supp } T$ is a subgrid in the grid-based case. A ratio set μ is a finite subset of $\mathfrak{G}^{\text{small}}$; $\mathfrak{J}^\mu$ is the group generated by μ . If $\mu = \{\mu_1, \dots, \mu_n\}$, then $\mathfrak{J}^\mu = \{\mu^{\mathbf{k}} : \mathbf{k} \in \mathbb{Z}^n\}$. If $\mathbf{m} \in \mathbb{Z}^n$, then $\mathfrak{J}^{\mu, \mathbf{m}} = \{\mu^{\mathbf{k}} : \mathbf{k} \in \mathbb{Z}^n, \mathbf{k} \geq \mathbf{m}\}$ is a grid. A grid-based transseries is supported by some grid. A subgrid is a subset of a grid.

For transseries A , we already use exponents A^n for multiplicative powers, and parentheses $A^{(n)}$ for derivatives. Therefore let us use square brackets $A^{[n]}$ for compositional powers. In particular, we will write $A^{[-1]}$ for the compositional inverse. Thus, for example, $\exp_n = \exp^{[n]} = \log^{[-n]}$.

In this paper we will be using some of the results on composition of transseries from [10] and [11], where the proofs are sometimes not as simple as in [9] (and not all of them are proved there).

1 Motivation for Fractional Iteration

Before we turn to transseries, let us consider fractional iteration in general. Given functions $T: X \rightarrow X$ and $\Phi: \mathbb{R} \times X \rightarrow X$, we say that Φ is a *real iteration group* for T iff

$$\Phi(s+t, x) = \Phi(s, \Phi(t, x)), \quad (1)$$

$$\Phi(0, x) = x, \quad (2)$$

$$\Phi(1, x) = T(x), \quad (3)$$

for all $s, t \in \mathbb{R}$ and $x \in X$.

Let us assume, say, that X is an interval $(a, b) \subseteq \mathbb{R}$, possibly $a = -\infty$ and/or $b = \infty$, and that Φ has as many derivatives as needed. If we start with (1), take the partial derivative with respect to t ,

$$\Phi_1(s+t, x) = \Phi_2(s, \Phi(t, x)) \Phi_1(t, x),$$

then substitute $t = 0$, we get

$$\Phi_1(s, x) = \Phi_2(s, x) \Phi_1(0, x). \quad (4)$$

We have written Φ_1 and Φ_2 for the two partial derivatives of Φ . Equation (4) is the one we will be using in Sections 2 and 3 below. Here is a proof showing how it works in the case of functions on an interval X .

Proposition 1.1. *Suppose $\Phi: \mathbb{R} \times X \rightarrow X$ satisfies (2) and*

$$\Phi_1(s, x) = \Phi_2(s, x) \beta(x) \quad (4')$$

with $\beta(x) > 0$. [Assume also $\infty = \int_{x_0}^b dy/\beta(y)$ and $\infty = \int_a^{x_0} dy/\beta(y)$ for $x_0 \in X = (a, b)$.] Then Φ satisfies (1) with $\beta(x) = \Phi_1(0, x)$.

Proof. Fix $x_0 \in X$. Let $\theta(t)$ be the solution of the ODE $\theta'(t) = \beta(\theta(t))$, $\theta(0) = x_0$. That is, $\theta(t) = x$ is defined implicitly by

$$\int_{x_0}^x \frac{dy}{\beta(y)} = t.$$

[In order to get all time t , we need $\infty = \int_{x_0}^b dy/\beta(y)$ and $\infty = \int_a^{x_0} dy/\beta(y)$.] Consider $F(u, v) = \Phi(u + v, \theta(u - v))$. Then

$$\begin{aligned} F_2(u, v) &= \Phi_1(u + v, \theta(u - v)) - \Phi_2(u + v, \theta(u - v)) \theta'(u - v) \\ &= \Phi_1(u + v, \theta(u - v)) - \Phi_2(u + v, \theta(u - v)) \beta(\theta(u - v)) = 0 \end{aligned}$$

by (4'). This means F is independent of v . So

$$\begin{aligned} \theta(t) &= \Phi(0, \theta(t)) = F(t/2, -t/2) = F(t/2, t/2) \\ &= \Phi(t, \theta(0)) = \Phi(t, x_0), \\ \Phi(s, \Phi(t, x_0)) &= \Phi(s, \theta(t)) = F((s + t)/2, (s - t)/2) \\ &= F((s + t)/2, (s + t)/2) = \Phi(s + t, \theta(0)) = \Phi(s + t, x_0). \end{aligned}$$

Differentiate $\Phi(t, x_0) = \theta(t)$ to get $\Phi_1(t, x_0) = \theta'(t) = \beta(\theta(t))$, then substitute $t = 0$ to get $\Phi_1(0, x_0) = \beta(\theta(0)) = \beta(x_0)$. $\square$

2 Three Examples

Power Series

We start with the classical case of power series. (A. Cayley 1860 [3]; A. Korkine 1882 [17].) We will think of formal power series (for $x \rightarrow \infty$), and not actual functions. Consider a series of the form

$$\begin{aligned} T(x) &= x \left(1 + \sum_{j=1}^{\infty} c_j x^{-j} \right) \\ &= x + \sum_{j=1}^{\infty} c_j x^{-j+1} = x + c_1 + c_2 x^{-1} + c_3 x^{-2} + \dots \end{aligned} \quad (5)$$

Such a series admits an iteration group of the same form. That is,

$$\Phi(s, x) = x \left(1 + \sum_{j=1}^{\infty} \alpha_j(s) x^{-j} \right). \quad (6)$$

In fact, $\alpha_j(s)$ is $sc_j + \{\text{polynomial in } s, c_1, c_2, \dots, c_{j-1} \text{ with rational coefficients, of degree } j-1 \text{ in } s\}$. The first few terms:

$$\begin{aligned}\Phi(s, x) = & x + sc_1 + sc_2x^{-1} + \left(sc_3 + \frac{s(1-s)}{2}c_1c_2 \right) x^{-2} \\ & + \left(sc_4 + \frac{s(1-s)}{2}(2c_1c_3 + c_2^2) + \frac{s(1-s)(1-2s)}{6}c_1^2c_2 \right) x^{-3} + \dots\end{aligned}$$

Theorem 2.1. Let $T(x)$ be the power series (5). Define $\alpha_j: \mathbb{R} \rightarrow \mathbb{R}$ recursively by

$$\begin{aligned}\alpha_1(s) &= sc_1, \\ \alpha_j(s) &= s \left(c_j - \int_0^1 \sum_{j_1+j_2=j} (-j_1+1)\alpha_{j_1}(u)\alpha'_{j_2}(0) du \right) \\ &\quad + \int_0^s \sum_{j_1+j_2=j} (-j_1+1)\alpha_{j_1}(u)\alpha'_{j_2}(0) du.\end{aligned}$$

Then the series Φ defined formally by (6) is a real iteration group for T .

Proof. Check (2), (3), (4). □

Remark 2.2. The formulas are obtained by plugging (6) into (4), equating coefficients, then integrating the resulting ODEs. Consequently, this is the *unique* solution of the form (6) with differentiable coefficients.

Remark 2.3. Of course there is a corresponding formulation for series of the form

$$T(z) = z \left(1 + \sum_{j=1}^{\infty} c_j z^j \right) = z + c_1 z^2 + c_2 z^3 + \dots$$

Then we get $\Phi(s, z) = z(1 + \sum_{j=1}^{\infty} \alpha_j(s)z^j)$, with

$$\begin{aligned}\alpha_1(s) &= sc_1, \\ \alpha_j(s) &= s \left(c_j - \int_0^1 \sum_{j_1+j_2=j} (j_1+1)\alpha_{j_1}(u)\alpha'_{j_2}(0) du \right) \\ &\quad + \int_0^s \sum_{j_1+j_2=j} (j_1+1)\alpha_{j_1}(u)\alpha'_{j_2}(0) du.\end{aligned}$$

The first few terms are:

$$\begin{aligned}\Phi(s, z) = & z + sc_1 z^2 + (sc_2 + s(s-1)c_1^2) z^3 \\ & + \left(sc_3 + \frac{5s(s-1)}{2}c_1c_2 + \frac{s(s-1)(2s-3)}{2}c_1^3 \right) z^4 + \dots\end{aligned}$$

Convergence

We considered here *formal* series. Even if (5) converges for all x (except 0), it need not follow that (6) converges. Indeed, one of the criticisms of Cayley [3] and Korkine [17] was that convergence was not proved. Baker [2] provides examples where a power series converges, but none of its non-integer iterates converges. Erdős & Jabotinsky [13] investigate the set of s for which the series converges.

Transseries, Height and Depth 0

Let $B \subseteq (0, \infty)$ be a well ordered set (under the usual order). The transseries

$$T(x) = x \left(1 + \sum_{b \in B} c_b x^{-b} \right) = x + \sum_{b \in B} c_b x^{-b+1} \quad (7)$$

is the next one we consider. In fact, the additive semigroup generated by B is again well ordered (Higman, see the proof in [10, Prop. 2.2]), so we will assume from the start that B is a semigroup. If B is finitely generated, then (7) is a grid-based transseries. But in general it is well-based. If B is finitely generated, then B has order type ω . But of course a well ordered B can have arbitrarily large countable ordinal as order type.

We claim T has a real iteration group supported by the same $1 - B$, where B is a well-ordered additive semigroup of positive reals.

Proposition 2.4. *Let $B \subseteq (0, \infty)$ be a semigroup. The transseries (7) has a real iteration group*

$$\Phi(s, x) = x \left(1 + \sum_{b \in B} \alpha_b(s) x^{-b} \right).$$

Proof. The only thing needed is that B is a well ordered semigroup. It follows that, for any given $b \in B$, there are just finitely many pairs $(b_1, b_2) \in B \times B$ with $b_1 + b_2 = b$ [9, Prop. 3.27]. Then define recursively:

$$\begin{aligned} f_b(s) &= \sum_{b_1 + b_2 = b} (-b_1 + 1) \alpha_{b_1}(s) \alpha'_{b_2}(0), \\ \alpha_b(s) &= s \left(c_b - \int_0^1 f_b(u) du \right) + \int_0^s f_b(u) du. \end{aligned}$$

For each b , both $f_b(s)$ and $\alpha_b(s)$ are polynomials (finitely many terms!) in s and the c_{b_1} [with $b_1 < b$ except for the term sc_b]. Check (2), (3), (4). $\square$

A Moderate Example

Now we consider another case. We single it out because it occurs frequently enough to make it useful to have the formulas displayed. Consider the transseries

$$T(x) = x \left(\sum_{k=0}^{\infty} \sum_{j=0}^{\infty} c_{j,k} x^{-j} e^{-kx} \right), \quad c_{0,0} = 1. \quad (8)$$

The set

$$B = \{ (j, k) : k \geq 0, j \geq 0, (j, k) \neq (0, 0) \} \quad (9)$$

is a semigroup under addition. The set $\{ x^{-j} e^{-kx} : (j, k) \in B \}$ is then a semigroup under multiplication. It is well ordered with order type ω^2 with respect to the converse of $\succ$.

Theorem 2.5. *Let B be as in (9). Then the transseries (8) admits a real iteration group supported by the same set $\{ x^{1-j} e^{-kx} : j, k \geq 0 \}$.*

Proof. Write

$$\Phi(s, x) = x \left(1 + \sum_{(j,k) \in B} \alpha_{j,k}(s) x^{-j} e^{-kx} \right).$$

(A) We first consider the case with $c_{1,0} = 0$. The coefficient functions $\alpha_{j,k}$ are defined recursively as follows. If $(j, k) \notin B$, then let $\alpha_{j,k}(s) = 0$. Let $(j, k) \in B$ and assume $\alpha_{j_1, k_1}(s)$ have already been defined for all (j_1, k_1) with either $k_1 < k$ or $\{k_1 = k \text{ and } j_1 < j\}$. Then let

$$f_{j,k}(s) = \sum [(-j_1 + 1)\alpha'_{j_2, k_2}(0) - k_1 \alpha'_{j_2+1, k_2}(0)] \alpha_{j_1, k_1}(s),$$

where the sum is over all j_1, j_2, k_1, k_2 with $j_1 + j_2 = j, k_1 + k_2 = k$. Check that all the terms in the sum involve α 's that have already been defined (or are multiplied by zero); this depends on $c_{1,0} = 0$, so $\alpha_{1,0}(s) = 0$. Define $F_{j,k}(s) = \int_0^s f_{j,k}(u) du$ and

$$\alpha_{j,k}(s) = (c_{j,k} - F_{j,k}(1))s + F_{j,k}(s).$$

Check (2), (3), (4).

(B) Now consider the case with $c_{1,0} \neq 0$. Write $c = c_{1,0}$. The coefficient functions $\alpha_{j,k}$ are defined recursively as follows. If $(j, k) \notin B$, then let $\alpha_{j,k}(s) = 0$. Let $(j, k) \in B$ and assume $\alpha_{j_1, k_1}(s)$ have already been defined for all (j_1, k_1) with either $k_1 < k$ or $\{k_1 = k \text{ and } j_1 < j\}$. Then let

$$f_{j,k}(s) = \sum [(-j_1 + 1)\alpha'_{j_2, k_2}(0) - k_1 \alpha'_{j_2+1, k_2}(0)] \alpha_{j_1, k_1}(s),$$

where the sum is over all j_1, j_2, k_1, k_2 with $j_1 + j_2 = j, k_1 + k_2 = k$. Omit the terms $\alpha'_{j,k}(0)$ and $-kc\alpha_{j,k}(s)$ and terms with a factor 0. Then all terms in the sum involve α 's already defined. Define $F_{j,k}(s) = \int_0^s e^{kc(u-s)} f_{j,k}(u) du$ and

$$\alpha_{j,k}(s) = \begin{cases} (c_{j,0} - F_{j,0}(1))s + F_{j,0}(s), & \text{if } k = 0, \\ (c_{j,k} - F_{j,k}(1)) \frac{1 - e^{-sck}}{1 - e^{-ck}} + F_{j,k}(s), & \text{if } k > 0. \end{cases}$$

Recall $c = c_{1,0}$. Check (2), (3), (4). □

In case (B)—the “moderate case”—the coefficients $\alpha_{j,k}(s)$ are not necessarily polynomials in s .

A Deep Example

Another simple example shows that real iteration group of that type need not always exist. Let $B \subseteq \mathbb{Z}^3$ be

$$B = \{(j, 0, 0) : j \geq 1\} \cup \{(j, k, 0) : k \geq 1\} \cup \{(j, k, l) : l \geq 1\}. \quad (10)$$

Then $\{x^{-j}e^{-kx}e^{-lx^2} : (j, k, l) \in B\}$ is a semigroup, but not well ordered.

I included some negative j and k so that the set of transseries of the form

$$x \left(1 + \sum_{(j,k,l) \in B} c_{jkl} x^{-j} e^{-kx} e^{-lx^2} \right)$$

(where of course each individual support is well ordered, not all of B) is closed under composition.

Proposition 2.6. *Let B' be a well ordered subset of (10). The transseries $T(x) = x(1 + x^{-1} + e^{-x^2})$ admits no real iteration group of the form*

$$\Phi(s, x) = x \left(1 + \sum_{(j,k,l) \in B'} \alpha_{jkl}(s) x^{-j} e^{-kx} e^{-lx^2} \right).$$

Proof. We may assume $B' = \{ (j, k, l) : \alpha_{jkl} \neq 0 \}$. As before, the first term beyond x can be computed as $\alpha_{100}(s) = s \cdot 1 = s$, so $\alpha'_{100}(s) = 1$. Now $\alpha_{001}(1) \neq 0$, so there is a least $(j, k, 1) \in B'$. But then by considering the coefficient of $x^{2-j} e^{-kx} e^{-x^2}$ in $\Phi_1(s, x) = \Phi_2(s, x) \Phi_1(0, x)$ we have

$$0 = (-2) \alpha_{jk1}(s) \alpha'_{100}(0),$$

so $\alpha_{jk1}(s) = 0$, a contradiction. $\square$

Remark 2.7. An alternate argument that there is no grid (or well ordered set) supporting a real iteration group for $T = x + 1 + xe^{-x^2}$. Compute

$$\begin{aligned} \text{supp } T^{[-1]} &= \{x \succ 1 \succ xe^{2x} e^{-x^2} \succ \dots\} \\ \text{supp } T^{[-2]} &= \{x \succ 1 \succ xe^{4x} e^{-x^2} \succ \dots\} \\ \text{supp } T^{[-3]} &= \{x \succ 1 \succ xe^{6x} e^{-x^2} \succ \dots\} \\ &\dots \\ \text{supp } T^{[-k]} &= \{x \succ 1 \succ xe^{2kx} e^{-x^2} \succ \dots\}, \quad k \in \mathbb{N}, k > 0. \end{aligned}$$

So there is no grid (and no well ordered set) containing all of these supports.

3 The Case of Common Support

Conjugation

The set $\mathcal{P}$ of large positive transseries is a group under composition ([10, Prop. 4.20], [11, Sec. 8]). In a group, we say U, V are **conjugate** if there exists S with $S^{[-1]} \circ U \circ S = V$. Then for all $k \in \mathbb{N}$ it follows that $S^{[-1]} \circ U^{[k]} \circ S = V^{[k]}$. If $\Phi(s, x)$ is an iteration group for $U(x)$, then $S^{[-1]}(\Phi(s, S(x)))$ is an iteration group for the conjugate $S^{[-1]}(U(S(x)))$. This can be used to reduce the question of fractional iteration for certain more general transseries to more restricted cases to be discussed here.

Puiseux series

For a Puiseux series of the form

$$T = \sum_{j=m}^{\infty} c_j x^{-j/k} \quad (11)$$

($m \in \mathbb{Z}, k \in \mathbb{N}, c_m > 0$), we can conjugate with $x^{1/k}$:

$$\begin{aligned} x^{1/k} \circ T \circ x^k &= \left(T(x^k) \right)^{1/k} = \left(\sum_{j=m}^{\infty} c_j x^{-j} \right)^{1/k} \\ &= c_m^{1/k} x^{-m/k} \left(1 + c_{m+1} x^{-1} + c_{m+2} x^{-2} + \dots \right)^{1/k} \\ &= c_m^{1/k} x^{-m/k} \left(1 + a_1 x^{-1} + a_2 x^{-2} + \dots \right). \end{aligned}$$

If $\text{dom } T = x$, then also $\text{dom}(x^{1/k} \circ T \circ x^k) = x$ and existence of a real iteration group is then clear from Proposition 5. (If $\text{dom } T \neq x$, keep reading.)

Exponentiality

Associated to a general $T \in \mathcal{P}$ is an integer p called the **exponentiality** of T [15, Ex. 4.10] [10, Prop. 4.5] such that for all large enough $k \in \mathbb{N}$ we have $\log_k \circ T \circ \exp_k \sim \exp_p$. Write $p = \text{expo } T$.

Now $\text{expo}(S \circ T) = \text{expo } S + \text{expo } T$, so no transseries with nonzero exponentiality can have a real iteration group of transseries. There is no transseries T with $T \circ T = e^x$. (But see [12].) The main question will be for exponentiality zero. If $\text{expo } T = 0$, then T is conjugate to some $S = \log_k \circ T \circ \exp_k$ such that $S \sim x$ and such that S is log-free [10, Prop. 4.8]. So we will deal with this case.

Shallow—Moderate—Deep

Now we turn to the general large positive log-free transseries with dominant term x . It admits a unique real iteration group with a common support in many cases (shallow and moderate), but not in many other cases (deep).

Definition 3.1. Consider log-free $T \sim x$. A real iteration group for T with **common support** is a real iteration group $\Phi(s, x)$ of the form

$$\Phi(s, x) = x \left(1 + \sum_{\mathfrak{g} \in \mathfrak{B}} \alpha_{\mathfrak{g}}(s) \mathfrak{g} \right) \quad (12)$$

for some subgrid ($\boxed{\mathbf{W}}$ or well ordered) $\mathfrak{B} \subseteq \mathfrak{G}$ (not depending on s) where coefficient functions $\alpha_{\mathfrak{g}}: \mathbb{R} \rightarrow \mathbb{R}$ are differentiable.

Write $T = x(1 + U)$, $U \prec 1$, $U \sim a\mathfrak{e}$, $a \in \mathbb{R}$, $a \neq 0$, $\mathfrak{e} \in \mathfrak{G}$, $\mathfrak{e} \prec 1$. As before, if there is a real iteration group Φ , it begins $\Phi(s, x) = x(1 + sa\mathfrak{e} + \dots)$. We may assume: if $\mathfrak{g} \in \mathfrak{B}$, then $\alpha_{\mathfrak{g}}(s) \neq 0$ for some s . So the greatest element of $\mathfrak{B}$ is $\mathfrak{e}$.

Write $A^\dagger = A'/A$ for the logarithmic derivative.

Definition 3.2. Let $T = x(1 + U)$, $U \prec 1$, $\text{mag } U = \mathfrak{e}$. Monomial $\mathfrak{e}$ is called the **first ratio** of T . We say that T is:

shallow iff $\mathfrak{g}^\dagger \prec 1/(x\mathfrak{e})$ for all $\mathfrak{g} \in \text{supp } U$;

moderate iff $\mathfrak{g}^\dagger \preccurlyeq 1/(x\mathfrak{e})$ for all $\mathfrak{g} \in \text{supp } U$ and $\mathfrak{g}^\dagger \succcurlyeq 1/(x\mathfrak{e})$ for at least one $\mathfrak{g} \in \text{supp } U$;

deep iff $\mathfrak{g}^\dagger \succcurlyeq 1/(x\mathfrak{e})$ for some $\mathfrak{g} \in \text{supp } U$.

purely deep iff $\mathfrak{g}^\dagger \succcurlyeq 1/(x\mathfrak{e})$ for all $\mathfrak{g} \in \text{supp } U$ except $\mathfrak{e}$.

Remark 3.3. $\boxed{\text{G}}$ It may be practical to check these definitions using a ratio set $\mu = \{\mu_1, \dots, \mu_n\}$. For example: Suppose $\text{supp } U \subseteq \mathfrak{J}^\mu$. By the group property (Lemma 3.14): if $\mu_i^\dagger \prec 1/(x\mathfrak{e})$ for $1 \leq i \leq n$, then T is shallow. This will be “if and only if” provided μ is chosen from the group generated by $\text{supp } U$, which can always be done.

Remark 3.4. The case of two terms, $T = x(1 + a\mathfrak{e})$ exactly, is shallow. Indeed, $\mathfrak{e} \prec 1$ and so $x\mathfrak{e} \prec x$, $x\mathfrak{e}' \prec 1$ and $\mathfrak{e}'/\mathfrak{e} \prec 1/(x\mathfrak{e})$.

Remark 3.5. For small monomials, the logarithmic derivative operation reverses the order: if $1 \succ \mathfrak{a} \succ \mathfrak{b}$, then $\mathfrak{a}^\dagger \preccurlyeq \mathfrak{b}^\dagger$ (Lemma 3.11(e)). And for $U \prec 1$ we have $U^\dagger \sim (\text{mag } U)^\dagger$ (Lemma 3.11(c)). So $T = x(1 + a\mathfrak{e} + V)$, $V \prec \mathfrak{e}$, is purely deep if and only if $V^\dagger \succcurlyeq 1/(x\mathfrak{e})$.

Remark 3.6. The condition $\mathfrak{g}^\dagger \prec 1/(x\mathfrak{e})$ says that $\mathfrak{g}$ is “not too small” in relation to $\mathfrak{e}$. (This is the reason for the terms “shallow” and “deep”.) If $\mathfrak{g} \prec 1$ then $\mathfrak{g} = e^{-L}$ with $L > 0$ purely large and

$$\begin{aligned} \mathfrak{g}^\dagger \prec \frac{1}{x\mathfrak{e}} &\iff L' \prec \frac{1}{x\mathfrak{e}} \\ &\iff L \prec \int \frac{1}{x\mathfrak{e}} \\ &\iff L < c \int \frac{1}{x\mathfrak{e}} \text{ for all real } c > 0 \\ &\iff e^L < \exp\left(c \int \frac{1}{x\mathfrak{e}}\right) \text{ for all real } c > 0 \\ &\iff \mathfrak{g} > \exp\left(-c \int \frac{1}{x\mathfrak{e}}\right) \text{ for all real } c > 0. \end{aligned}$$

So the set $\mathfrak{A} = \{\mathfrak{g} \in \mathfrak{G} : \mathfrak{g} \preccurlyeq \mathfrak{e}, \mathfrak{g}^\dagger \prec 1/(x\mathfrak{e})\}$ is an **interval** in $\mathfrak{G}$. The large end of the interval is the first ratio $\mathfrak{e}$, the small end of the interval is the gap in $\mathfrak{G}$ just above all the values $\exp(-c \int (1/x\mathfrak{e}))$, $c \in \mathbb{R}$, $c > 0$. If we write $\exp(-\bar{0} \int (1/x\mathfrak{e}))$ for that gap, then

$$\mathfrak{A} =] \exp(-\bar{0} \int (1/x\mathfrak{e})), \mathfrak{e}].$$

I will call this the **shallow interval** below $\mathfrak{e}$. Van der Hoeven devotes a chapter [15, Chap. 9] to gaps (cuts) in the transline. In his classification [15, Prop. 9.15],

$$\exp(-\bar{0} \int (1/x\mathfrak{e})) = e^{-e^A - e^{\bar{0}}},$$

where $\text{mag } \int (1/(x\mathfrak{e})) = e^A$.

Similarly, the set $\mathfrak{A} = \{\mathfrak{g} \in \mathfrak{G} : \mathfrak{g} \preccurlyeq \mathfrak{e}, \mathfrak{g}^\dagger \preccurlyeq 1/(x\mathfrak{e})\}$ is an interval in $\mathfrak{G}$. The large end of the interval is the first ratio $\mathfrak{e}$, the small end of the interval is the gap in $\mathfrak{G}$ just

below all the values $\exp(-c \int(1/x\epsilon))$, $c \in \mathbb{R}$, $c > 0$. If we write $\exp(-\overline{\infty} \int(1/x\epsilon))$ for that gap, then

$$\mathfrak{A} =] \exp(-\overline{\infty} \int(1/x\epsilon)), \epsilon].$$

I will call this the **moderate interval** below ϵ . In van der Hoeven's classification,

$$\exp(-\overline{\infty} \int(1/x\epsilon)) = e^{-e^A + e^{\overline{\sigma}}},$$

where $\text{mag} \int(1/(x\epsilon)) = e^A$.

The Examples

Let us examine where the examples done above fit in the shallow/deep classification. If $\epsilon = x^{-1}$, then $1/(x\epsilon) = 1$, $\int(1/(x\epsilon)) = x$, $\exp(-c \int(1/(x\epsilon))) = e^{-cx}$. The small end of the shallow interval is $\exp(-\overline{0}x)$. In a power series, every monomial $x^{-j} \succ e^{-\overline{0}x}$ is inside the shallow interval, so a power series is shallow.

In example (8), we saw two cases. In case $c_{1,0} \neq 0$, then $\epsilon = x^{-1}$ so again the small end of the shallow interval is $\exp(-\overline{0}x)$. But the monomial $x^{-j}e^{-kx} \prec \exp(-\overline{0}x)$ if $k > 0$, and is thus outside the shallow interval, so this is not shallow. The small end of the moderate interval is $\exp(-\overline{\infty}x)$, and all monomials $x^{-j}e^{-kx} \succ \exp(-\overline{\infty}x)$ are inside the moderate interval, so this is the moderate case.

The other case is $c_{1,0} = 0$. Then ϵ is x^{-2} (or smaller). If $\epsilon = x^{-2}$, then $1/(x\epsilon) = x$, $\exp(-c \int(1/(x\epsilon))) = e^{-(c/2)x^2}$. The small end of the shallow interval is $\exp(-\overline{0}x^2)$. All monomials $x^{-j}e^{-kx} \succ \exp(-\overline{0}x^2)$ are inside the shallow interval, so this is the shallow case.

Finally consider the example $T = x(1 + x^{-1} + e^{-x^2})$ of Proposition 2.6. Since $\epsilon = x^{-1}$, the small end of the moderate interval was computed as $\exp(-\overline{\infty}x)$. The monomial $e^{-x^2} \prec \exp(-\overline{\infty}x)$ is outside of that, so T is deep.

Proofs

Proofs will follow the examples done above. These proofs use some technical lemmas on logarithmic derivatives, grids, and well ordered sets; they are found after the main results, starting with Lemma 3.11.

Theorem 3.7. *If (log-free) $T \sim x$ is shallow, then T admits a real iteration group with common support where all coefficient functions are polynomials.*

Proof. The proof is as in Proposition 2.4 above. Here are the details. Write $T = x(1 + U)$, $U \sim a\epsilon$, $\epsilon \prec 1$. Begin with the subgrid ($\boxed{\mathbb{W}}$ well ordered) $\text{supp } U$ which is contained in $\{\mathfrak{g} \in \mathfrak{G} : \mathfrak{g} \succ \epsilon, \mathfrak{g}^\dagger \prec 1/(x\epsilon)\}$. Let $\mathfrak{B} \supseteq \text{supp } U$ be the least set such that if $\mathfrak{g}_1, \mathfrak{g}_2 \in \mathfrak{B}$, then $\text{supp}((x\mathfrak{g}_1)'\mathfrak{g}_2) \subseteq \mathfrak{B}$. By Lemma 3.22, $\mathfrak{B}$ is a subgrid ($\boxed{\mathbb{W}}$ by Lemma 3.23, $\mathfrak{B}$ is well ordered) and $\mathfrak{B} \subseteq \{\mathfrak{g} \in \mathfrak{G} : \mathfrak{g} \succ \epsilon, \mathfrak{g}^\dagger \prec 1/(x\epsilon)\}$.

Write

$$T(x) = x \left(1 + \sum_{\mathfrak{g} \in \mathfrak{B}} c_{\mathfrak{g}} \mathfrak{g} \right)$$

$$\begin{aligned}
\Phi(s, x) &= x \left(1 + \sum_{\mathfrak{g} \in \mathfrak{B}} \alpha_{\mathfrak{g}}(s) \mathfrak{g} \right) \\
\Phi_1(s, x) &= x \sum_{\mathfrak{g} \in \mathfrak{B}} \alpha'_{\mathfrak{g}}(s) \mathfrak{g} \\
\Phi_1(0, x) &= x \sum_{\mathfrak{g} \in \mathfrak{B}} \alpha'_{\mathfrak{g}}(0) \mathfrak{g} \\
\Phi_2(s, x) &= 1 + \sum_{\mathfrak{g} \in \mathfrak{B}} \alpha_{\mathfrak{g}}(s) (x\mathfrak{g})' \\
\Phi_2(s, x) \Phi_1(0, x) &= x \sum_{\mathfrak{g} \in \mathfrak{B}} \alpha'_{\mathfrak{g}}(0) \mathfrak{g} + x \sum_{\mathfrak{g}_1, \mathfrak{g}_2 \in \mathfrak{B}} \alpha_{\mathfrak{g}_1}(s) \alpha'_{\mathfrak{g}_2}(0) (x\mathfrak{g}_1)' \mathfrak{g}_2.
\end{aligned} \tag{13}$$

Now fix a monomial $\mathfrak{g} \in \mathfrak{B}$. Assume $\alpha_{\mathfrak{g}_1}$ has been defined for all $\mathfrak{g}_1 \succ \mathfrak{g}$. Consideration of the coefficient of $x\mathfrak{g}$ in (4) gives us an equation

$$\alpha'_{\mathfrak{g}}(s) = \alpha'_{\mathfrak{g}}(0) + f_{\mathfrak{g}}(s), \tag{14}$$

where $f_{\mathfrak{g}}(s)$ is a (real) linear combination of terms $\alpha_{\mathfrak{g}_1}(s) \alpha'_{\mathfrak{g}_2}(0)$ where $\mathfrak{g}_1, \mathfrak{g}_2 \in \mathfrak{B}$ satisfy $\mathfrak{g} \in \text{supp}((x\mathfrak{g}_1)'\mathfrak{g}_2)$. By Lemma 3.19 ([W] Lemma 3.20), for a given value of $\mathfrak{g}$, there are only finitely many pairs $\mathfrak{g}_1, \mathfrak{g}_2$ involved.

Now I claim these all have $\mathfrak{g}_1 \succ \mathfrak{g}$ and $\mathfrak{g}_2 \succ \mathfrak{g}$. Indeed, since $\mathfrak{g} \in \text{supp}((x\mathfrak{g}_1)'\mathfrak{g}_2)$ and $(x\mathfrak{g}_1)' = \mathfrak{g}_1 + x\mathfrak{g}_1'$, we have either $\mathfrak{g} \preceq \mathfrak{g}_1\mathfrak{g}_2$ or $\mathfrak{g} \preceq x\mathfrak{g}_1'\mathfrak{g}_2$. Take two cases:

- (a) $\mathfrak{g} \preceq \mathfrak{g}_1\mathfrak{g}_2$: Now $\mathfrak{g}_1 \preceq \mathfrak{e}$, so $\mathfrak{g} \preceq \mathfrak{e}\mathfrak{g}_2 \prec \mathfrak{g}_2$. And $\mathfrak{g}_2 \preceq \mathfrak{e}$, so $\mathfrak{g} \preceq \mathfrak{e}\mathfrak{g}_1 \prec \mathfrak{g}_1$.
- (b) $\mathfrak{g} \preceq x\mathfrak{g}_1'\mathfrak{g}_2$: Now $\mathfrak{g}_1' \prec \mathfrak{g}_1/(x\mathfrak{e})$ so $\mathfrak{g} \preceq x\mathfrak{g}_1'\mathfrak{g}_2 \prec \mathfrak{g}_1\mathfrak{g}_2/\mathfrak{e}$. But $\mathfrak{g}_1 \preceq \mathfrak{e}$, so $\mathfrak{g} \prec \mathfrak{g}_1$. And $\mathfrak{g}_2 \preceq \mathfrak{e}$, so $\mathfrak{g} \prec \mathfrak{g}_2$.

Thus we may use equations (14) to recursively define $\alpha_{\mathfrak{g}}(s)$. Indeed, solving the differential equation, we get $\alpha_{\mathfrak{g}}(s) = \int_0^s f_{\mathfrak{g}}(u) du + s\alpha'_{\mathfrak{g}}(0) + C$; but $\alpha_{\mathfrak{g}}(0) = 0$ and $\alpha_{\mathfrak{g}}(1) = c_{\mathfrak{g}}$, so

$$\alpha_{\mathfrak{g}}(s) = s \left(c_{\mathfrak{g}} - \int_0^1 f_{\mathfrak{g}}(u) du \right) + \int_0^s f_{\mathfrak{g}}(u) du. \tag{15}$$

In particular, the recursion begins with $\mathfrak{g} = \mathfrak{e}$ where $f_{\mathfrak{e}}(s) = 0$ and $\alpha_{\mathfrak{e}}(s) = sc_{\mathfrak{e}} = sa$. Also (by induction) $\alpha_{\mathfrak{g}}(s)$ and $f_{\mathfrak{g}}(s)$ are polynomials in s . $\square$

Theorem 3.8. *If (log-free) $T \sim x$ is moderate, then T admits a real iteration group with common support. The coefficient functions are entire; we cannot conclude the coefficient functions are polynomials.*

Proof. The proof is as in Theorem 2.5 above. Here are the details. Write $T = x(1+U)$, $U \sim a\mathfrak{e}$, $\mathfrak{e} \prec 1$. Begin with the subgrid ([W] well ordered) $\text{supp } U$ which is contained in $\{\mathfrak{g} \in \mathfrak{G} : \mathfrak{g} \succ \mathfrak{e}, \mathfrak{g}^\dagger \preceq 1/(x\mathfrak{e})\}$. Let $\mathfrak{B} \supseteq \text{supp } U$ be the least set such that if $\mathfrak{g}_1, \mathfrak{g}_2 \in \mathfrak{B}$, then $\text{supp}((x\mathfrak{g}_1)'\mathfrak{g}_2) \subseteq \mathfrak{B}$. By Lemma 3.22, $\mathfrak{B}$ is a subgrid ([W] by Lemma 3.23, $\mathfrak{B}$ is well ordered) and $\mathfrak{B} \subseteq \{\mathfrak{g} \in \mathfrak{G} : \mathfrak{g} \succ \mathfrak{e}, \mathfrak{g}^\dagger \preceq 1/(x\mathfrak{e})\}$.

Write $T(x) = x(1 + \sum_{\mathfrak{g} \in \mathfrak{B}} c_{\mathfrak{g}}\mathfrak{g})$ and $\Phi(s, x) = x(1 + \sum_{\mathfrak{g} \in \mathfrak{B}} \alpha_{\mathfrak{g}}(s)\mathfrak{g})$. Compute the derivatives as in (13).

Now fix a monomial $\mathfrak{g} \in \mathfrak{B}$. Assume $\alpha_{\mathfrak{g}_1}$ has been defined for all $\mathfrak{g}_1 \succ \mathfrak{g}$. There are two cases. If $\mathfrak{g}^\dagger \prec 1/(x\mathfrak{e})$, then the argument proceeds as before, and we get (15). Now

assume $\mathfrak{g}^\dagger \asymp 1/(x\mathfrak{e})$, say $\mathfrak{g}^\dagger \sim b/(x\mathfrak{e})$, $b \in \mathbb{R}$, $b \neq 0$. Consideration of the coefficient of $x\mathfrak{g}$ in (4) gives us an equation

$$\alpha'_\mathfrak{g}(s) = \alpha'_\mathfrak{g}(0) + f_\mathfrak{g}(s) + b\alpha_\mathfrak{g}(s)\alpha'_\mathfrak{e}(0), \quad (16)$$

where $f_\mathfrak{g}(s)$ is a (real) linear combination of terms $\alpha_{\mathfrak{g}_1}(s)\alpha'_{\mathfrak{g}_2}(0)$ where $\mathfrak{g}_1, \mathfrak{g}_2 \in \mathfrak{B}$ satisfy $\mathfrak{g} \in \text{supp}((x\mathfrak{g}_1)'\mathfrak{g}_2)$. The term with $\mathfrak{g}_1 = \mathfrak{g}$, $\mathfrak{g}_2 = \mathfrak{e}$, namely $b\alpha_\mathfrak{g}(s)\alpha'_\mathfrak{e}(0)$, is not included in $f_\mathfrak{g}$ but written separately. By Lemma 3.19 ([W] Lemma 3.20), for a given value of $\mathfrak{g}$, there are only finitely many pairs $\mathfrak{g}_1, \mathfrak{g}_2$ involved.

But I claim these all have $\mathfrak{g}_1 \succ \mathfrak{g}$ and $\mathfrak{g}_2 \succ \mathfrak{g}$ except for the case $\mathfrak{g}_1 = \mathfrak{g}, \mathfrak{g}_2 = \mathfrak{e}$. Indeed, since $\mathfrak{g} \in \text{supp}((x\mathfrak{g}_1)'\mathfrak{g}_2)$ and $(x\mathfrak{g}_1)' = \mathfrak{g}_1 + x\mathfrak{g}'_1$, we have either $\mathfrak{g} \preccurlyeq \mathfrak{g}_1\mathfrak{g}_2$ or $\mathfrak{g} \preccurlyeq x\mathfrak{g}'_1\mathfrak{g}_2$. Take two cases:

- (a) $\mathfrak{g} \preccurlyeq \mathfrak{g}_1\mathfrak{g}_2$: Now $\mathfrak{g}_1 \preccurlyeq \mathfrak{e}$, so $\mathfrak{g} \preccurlyeq \mathfrak{e}\mathfrak{g}_2 \prec \mathfrak{g}_2$. And $\mathfrak{g}_2 \preccurlyeq \mathfrak{e}$, so $\mathfrak{g} \preccurlyeq \mathfrak{e}\mathfrak{g}_1 \prec \mathfrak{g}_1$.
- (b) $\mathfrak{g} \preccurlyeq x\mathfrak{g}'_1\mathfrak{g}_2$: Now $\mathfrak{g}'_1 \preccurlyeq \mathfrak{g}_1/(x\mathfrak{e})$ so $\mathfrak{g} \preccurlyeq x\mathfrak{g}'_1\mathfrak{g}_2 \preccurlyeq \mathfrak{g}_1\mathfrak{g}_2/\mathfrak{e}$, with $\asymp$ only if $\mathfrak{g} \asymp x\mathfrak{g}'_1\mathfrak{g}_2$ and $\mathfrak{g}_1^\dagger = 1/(x\mathfrak{e})$. But $\mathfrak{g}_1 \preccurlyeq \mathfrak{e}$, so $\mathfrak{g} \preccurlyeq \mathfrak{g}_1$, and $\asymp$ would mean $\mathfrak{g}_1 = \mathfrak{e}$ and $\mathfrak{g}_1^\dagger = 1/(x\mathfrak{e})$ so $\mathfrak{e}^\dagger \asymp 1/(x\mathfrak{e})$, which is false; thus $\mathfrak{g} \prec \mathfrak{g}_1$. And $\mathfrak{g}_2 \preccurlyeq \mathfrak{e}$, so $\mathfrak{g} \preccurlyeq \mathfrak{g}_2$. This time $\asymp$ means $\mathfrak{g}_2 = \mathfrak{e}$, $\mathfrak{g}_1 = \mathfrak{g}$, $\mathfrak{g}^\dagger \asymp 1/(x\mathfrak{e})$.

Thus we may use equations (16) to recursively define $\alpha_\mathfrak{g}(s)$. Indeed, solving the differential equation, we get

$$\alpha_\mathfrak{g}(s) = \frac{e^{abs} - 1}{e^{ab} - 1} \left(c_\mathfrak{g} - \int_0^1 e^{ab(1-u)} f_\mathfrak{g}(u) du \right) + \int_0^s e^{ab(s-u)} f_\mathfrak{g}(u) du \quad (17)$$

if $\mathfrak{g}^\dagger \sim b/(x\mathfrak{e})$. □

Theorem 3.9. *If (log-free) $T \sim x$ is deep, then it does not admit a real iteration group of the form (12).*

Proof. The proof is as in Proposition 2.6 above. Here are the details. Write $T = x(1+U)$, $U \sim a\mathfrak{e}$, $\mathfrak{e} \prec 1$. Suppose

$$\Phi(s, x) = x \left(1 + \sum_{\mathfrak{g} \in \mathfrak{B}} \alpha_\mathfrak{g}(s)\mathfrak{g} \right)$$

satisfies (4), (2), $\Phi(1, x) = T(x)$, and $\mathfrak{B}$ is a subgrid ([W] well ordered). We may assume $\alpha_\mathfrak{g} \neq 0$ for all $\mathfrak{g} \in \mathfrak{B}$. Write

$$\mathfrak{B}_1 = \left\{ \mathfrak{g} \in \mathfrak{B} : \mathfrak{g}^\dagger \preccurlyeq 1/(x\mathfrak{e}) \right\}, \quad \mathfrak{B}_2 = \left\{ \mathfrak{g} \in \mathfrak{B} : \mathfrak{g}^\dagger \succ 1/(x\mathfrak{e}) \right\}$$

(the moderate and deep portions of $\mathfrak{B}$). As before, $\max \mathfrak{B} = \mathfrak{e}$ and $\alpha_\mathfrak{e}(s) = as$, $\alpha'_\mathfrak{e}(0) = a$, $a \neq 0$.

Let $\mathfrak{m} = \max \mathfrak{B}_2$. So $\mathfrak{m}^\dagger \succ 1/(x\mathfrak{e})$. Write $\mathfrak{m}^\dagger \sim -b\mathfrak{p}/(x\mathfrak{e})$, $b \in \mathbb{R}$, $b > 0$, $\mathfrak{p} \in \mathfrak{G}$, $\mathfrak{p} \succ 1$. ($\mathfrak{m}$ is small and positive, so $\mathfrak{m}'$ is negative.) In (4) we will consider the coefficient of $x\mathfrak{p}\mathfrak{m} = x \text{mag}((x\mathfrak{m})'\mathfrak{e})$. By Lemma 3.11(g), $((x\mathfrak{m})')^\dagger \sim \mathfrak{m}^\dagger \succ 1/(x\mathfrak{e})$; by Remark 3.4, $\mathfrak{e}^\dagger \prec 1/(x\mathfrak{e})$; so $(\mathfrak{p}\mathfrak{m})^\dagger = ((x\mathfrak{m})')^\dagger + \mathfrak{e}^\dagger \sim ((x\mathfrak{m})')^\dagger \succ 1/(x\mathfrak{e})$ and $\mathfrak{p}\mathfrak{m} \notin \mathfrak{B}_1$. Also $\mathfrak{p}\mathfrak{m} \succ \mathfrak{m}$, so $\mathfrak{p}\mathfrak{m} \notin \mathfrak{B}_2$. Thus $\mathfrak{p}\mathfrak{m} \notin \mathfrak{B}$ so $\alpha_{\mathfrak{p}\mathfrak{m}} = 0$.

I claim the only $\mathfrak{g}_1, \mathfrak{g}_2 \in \mathfrak{B}$ with $\mathfrak{p}\mathfrak{m} \in \text{supp}((x\mathfrak{g}_1)'\mathfrak{g}_2)$ are $\mathfrak{g}_1 = \mathfrak{m}$, $\mathfrak{g}_2 = \mathfrak{e}$. To prove this, consider these cases:

- (a) $\mathfrak{g}_1 \succ \mathfrak{m}, \mathfrak{g}_2 \succ \mathfrak{m}$. By Lemma 3.14(b), $\text{supp}((x\mathfrak{g}_1)') \subseteq \mathfrak{B}_1$ and by Lemma 3.14(a), $\text{supp}((x\mathfrak{g}_1)'\mathfrak{g}_2) \subseteq \mathfrak{B}_1$. So $\mathfrak{pm} \notin \text{supp}((x\mathfrak{g}_1)'\mathfrak{g}_2)$.
- (b) $\mathfrak{g}_1 = \mathfrak{m}, \mathfrak{g}_2 = \mathfrak{e}$. In this case $\mathfrak{pm} = \text{mag}((x\mathfrak{g}_1)'\mathfrak{g}_2)$, so $\mathfrak{pm} \in \text{supp}((x\mathfrak{g}_1)'\mathfrak{g}_2)$.
- (c) $\mathfrak{g}_1 = \mathfrak{m}, \mathfrak{g}_2 \prec \mathfrak{e}$. Then $(x\mathfrak{g}_1)'\mathfrak{g}_2 \prec (x\mathfrak{m})'\mathfrak{e} \asymp \mathfrak{pm}$, so $\mathfrak{pm} \notin \text{supp}((x\mathfrak{g}_1)'\mathfrak{g}_2)$.
- (d) $\mathfrak{g}_1 \prec \mathfrak{m}$. Then $(x\mathfrak{g}_1)'\mathfrak{g}_2 \prec (x\mathfrak{m})'\mathfrak{e} \asymp \mathfrak{pm}$, so $\mathfrak{pm} \notin \text{supp}((x\mathfrak{g}_1)'\mathfrak{g}_2)$.
- (e) $\mathfrak{g}_2 \asymp \mathfrak{m}$. Then by Remark 3.4, $(x\mathfrak{g}_1)'\mathfrak{g}_2 \asymp (x\mathfrak{e})'\mathfrak{m} \prec \mathfrak{m} \prec \mathfrak{pm}$, so $\mathfrak{pm} \notin \text{supp}((x\mathfrak{g}_1)'\mathfrak{g}_2)$.

So consideration of the coefficient of $x\mathfrak{pm}$ in (4) yields:

$$0 = -b\alpha_{\mathfrak{m}}(s)\alpha'_{\mathfrak{e}}(0) = -ab\alpha_{\mathfrak{m}}(s),$$

where $-ab \neq 0$, so $\alpha_{\mathfrak{m}} = 0$, a contradiction. $\square$

Remark 3.10. If $T \sim x$, then a calculation shows that T is shallow, moderate, or deep if and only if $\log \circ T \circ \exp$ is shallow, moderate, or deep, respectively. This invariance of the classification will show that the three theorems above are correct for all $T \in \mathbb{T}$ with $T \sim x$, even if not log-free. For example, $T = x + \log x$ is shallow, since log-free

$$\log \circ T \circ \exp = x + \sum_{j=1}^{\infty} \frac{(-1)^{j+1}}{j} x^j e^{-jx}$$

is shallow.

Recall that if T is large and positive and has exponentiability 0, then for some k we have $\log^{[k]} \circ T \circ \exp^{[k]} \sim x$. We might try to use the same principle to extend the definitions of shallow, moderate, and deep to such transseries T even if $T \not\sim x$. Define T **shallow** provided $\log^{[k]} \circ T \circ \exp^{[k]}$ is shallow for some k ; similarly for moderate and deep. Examples: $T = x \log x$ is shallow, since $\log \circ T \circ \exp = x + \log x$ is shallow. The finite power series $U = 2x - 2/x$ is moderate, since

$$\log \circ U \circ \exp = x + \log 2 - \sum_{j=1}^{\infty} \frac{e^{-2jx}}{j}$$

is moderate. And $V = 2x - 2e^{-x}$ is deep.

But the usefulness of this extension is not entirely clear, since it may produce a family $\Phi(s, x)$ without common support. Example: For $T = x \log x$ we compute $S = \log \circ T \circ \exp = x + \log x$. So we get a real iteration group for S of the form $\Psi(s, x) = x + s \log x + o(1)$, and then a real iteration group for T of the form $\Phi(s, x) = \exp(\Psi(s, \log x)) = x(\log x)^s + \dots$. These are not all supported by a common subgrid or even a common well ordered set. When the support depends on the parameter s , it may no longer make sense to require the coefficients be differentiable.

Example $x^2 + c$ is also deep. It is discussed in Section 6, below. The figure there illustrates supports of iterates $M^{[s]}$ that vary with s .

Technical Lemmas

Lemma 3.11. *Let $\mathfrak{a}, \mathfrak{b} \in \mathfrak{G}$ and $A, B \in \mathbb{T}$. (a) If $\mathfrak{a} = e^A$, with $A \neq 0$ purely large, then $\mathfrak{a}^\dagger = A'$. (b) $(AB)^\dagger = A^\dagger + B^\dagger$. (c) If $A = a\mathfrak{g}(1 + U)$, $\mathfrak{g} \neq 1$, $U \prec 1$, then $A^\dagger \sim \mathfrak{g}^\dagger$. (d) If $1 \prec \mathfrak{a} \leq \mathfrak{b}$, then $\mathfrak{a}^\dagger \leq \mathfrak{b}^\dagger$. (d') If $1 \prec A \leq B$, then $A^\dagger \leq B^\dagger$. (e) If $1 \succ \mathfrak{a} \geq \mathfrak{b}$, then $\mathfrak{a}^\dagger \leq \mathfrak{b}^\dagger$. (e') If $1 \succ A \geq B$, then $A^\dagger \leq B^\dagger$. (f) If $\mathfrak{b} \prec \mathfrak{a} \prec 1/\mathfrak{b}$, then $\mathfrak{a}^\dagger \leq \mathfrak{b}^\dagger$. (g) If $\mathfrak{b} \neq 1$ is log-free and $\mathfrak{n} \in \text{supp}((x\mathfrak{b})')$, then $\mathfrak{n}^\dagger \sim \mathfrak{b}^\dagger$.*

Proof. (a) $\mathfrak{a}' = A'e^A$ so $\mathfrak{a}^\dagger = \mathfrak{a}'/\mathfrak{a} = A'$.

(b) Product rule.

(c) Since $\mathfrak{g} = e^L$, L purely large, note $\mathfrak{g}^\dagger = L'$. Also $U \prec 1$ so $(1+U)^\dagger = U'/(1+U) \sim U'$. Now $L \succ 1 \succ U$, so $L' \succ U'$ and $\mathfrak{g}^\dagger \succ (1+U)^\dagger$. So $A^\dagger = \mathfrak{a}^\dagger + \mathfrak{g}^\dagger + (1+U)^\dagger \sim \mathfrak{g}^\dagger$.

(d) Write $\mathfrak{a} = e^A$, $\mathfrak{b} = e^B$, with A, B purely large. Now $0 < A \leq B$, so $A \preceq B$. Also $B \not\prec 1$, so $A' \preceq B'$, that is, $\mathfrak{a}^\dagger \preceq \mathfrak{b}^\dagger$. (d') follows from (c).

(e) Write $\mathfrak{a} = e^A$, $\mathfrak{b} = e^B$, with A, B purely large. Now $0 > A \geq B$, so $A \preceq B$. Also $B \not\prec 1$, so $A' \preceq B'$, that is, $\mathfrak{a}^\dagger \preceq \mathfrak{b}^\dagger$. (e') follows from (c).

(f) If $\mathfrak{a} = 1$, then $\mathfrak{a}^\dagger = 0$. If $\mathfrak{a} \prec 1$, then apply (e). If $\mathfrak{a} \succ 1$, note that $\mathfrak{b}^\dagger = -(1/\mathfrak{b})^\dagger \asymp (1/\mathfrak{b})^\dagger$, then apply (d).

(g) Case 1: $\mathfrak{b} = x^b$, $b \in \mathbb{R}$, $b \neq 0$. Then $x\mathfrak{b} = x^{b+1}$, $(x\mathfrak{b})' = (b+1)x^b \sim \mathfrak{b}$. So $((x\mathfrak{b})')^\dagger \sim \mathfrak{b}^\dagger$. Case 2: $\mathfrak{b} = x^b e^L$, $L \neq 0$ has height $n \geq 0$. Assume $\mathfrak{b} \prec 1$, the case $\mathfrak{b} \succ 1$ is similar. Then $x\mathfrak{b} = x^{b+1} e^L$, $(x\mathfrak{b})' = ((b+1) + xL')x^b e^L = A\mathfrak{b}$ where $A = (b+1) + xL'$ has height n . Now $\mathfrak{n} \in \text{supp}((x\mathfrak{b})')$, so $\mathfrak{n} = \mathfrak{a}\mathfrak{b}$ where $\mathfrak{a} \in \text{supp} A$. But $\mathfrak{a}$ has height n , so by “height wins” [9, Prop. 3.72], we have $\mathfrak{b} \prec \mathfrak{a} \prec 1/\mathfrak{b}$. In the proofs of (d) and (e), if it is a case of “height wins” then we get strict inequality $A \prec B$ and $\mathfrak{a}^\dagger \prec \mathfrak{b}^\dagger$. So here $\mathfrak{a}^\dagger \prec \mathfrak{b}^\dagger$. Therefore $\mathfrak{n}^\dagger = (\mathfrak{a}\mathfrak{b})^\dagger = \mathfrak{a}^\dagger + \mathfrak{b}^\dagger \sim \mathfrak{b}^\dagger$. $\square$

Lemma 3.12. *Let $T = x(1 + a\epsilon + o(\epsilon))$ with $\epsilon \in \mathfrak{G}$, $\epsilon \prec 1$, $a \in \mathbb{R}$, $a > 0$. Let $\mathfrak{g} \in \mathfrak{G}$, $\mathfrak{g} \prec 1$. Then (i) $\epsilon(T) \sim \epsilon$; (ii) if $\mathfrak{g} \prec 1/(x\epsilon)$ then $\mathfrak{g}(T) \sim \mathfrak{g}$; (iii) if $\mathfrak{g} \sim -b/(x\epsilon)$, $b \in \mathbb{R}$ then $\mathfrak{g}(T) \sim e^{-ab}\mathfrak{g}$; (iv) if $\mathfrak{g} \succ 1/(x\epsilon)$, then $\mathfrak{g}(T) \prec \mathfrak{g}$.*

Proof. (i) As in Remark 3.4, $\epsilon' \prec 1/x$. Then

$$\epsilon(T) - \epsilon = \int_x^T \epsilon' \prec \int_x^T \frac{1}{x} = \log \frac{T}{x} = \log(1 + a\epsilon + o(\epsilon)) \sim a\epsilon \asymp \epsilon,$$

so $\epsilon(T) \sim \epsilon$.

(ii) Assume $\mathfrak{g}^\dagger \prec 1/(x\epsilon)$. Then

$$\log \frac{\mathfrak{g}(T)}{\mathfrak{g}} = \int_x^T \mathfrak{g}^\dagger \prec \int_x^T \frac{1}{x\epsilon}.$$

By [10, Prop. 4.24], the value of this integral is between

$$\frac{T-x}{x\epsilon} \sim \frac{ax\epsilon}{x\epsilon} = a \quad \text{and} \quad \frac{T-x}{T\epsilon(T)} \sim \frac{ax\epsilon}{x\epsilon} = a.$$

Thus $\log(\mathfrak{g}(T)/\mathfrak{g}) \prec 1$ so $\mathfrak{g}(T)/\mathfrak{g} \sim 1$.

(iii) Assume $\mathfrak{g}^\dagger \sim -b/(x\epsilon)$. Then

$$\log \frac{\mathfrak{g}(T)}{\mathfrak{g}} = \int_x^T \mathfrak{g}^\dagger \sim \int_x^T \frac{-b}{x\epsilon} \sim -ab,$$

where the integral was estimated in the same way as in (ii). Therefore $\mathfrak{g}(T) \sim e^{-ab}\mathfrak{g}$.

(iv) Assume $\mathfrak{g}^\dagger \succ 1/(x\epsilon)$. Then

$$\log \frac{\mathfrak{g}(T)}{\mathfrak{g}(x)} = \int_x^T \mathfrak{g}^\dagger \succ \int_x^T \frac{1}{x\epsilon} \sim a.$$

But $\log(\mathfrak{g}(T)/\mathfrak{g}(x)) < 0$ so $\mathfrak{g}(T)/\mathfrak{g}(x) \prec 1$ and $\mathfrak{g}(T) \prec \mathfrak{g}$. $\square$

For completeness, we note the following analogous version for $T < x$.

Lemma 3.13. *Let $T = x(1 - a\epsilon + o(\epsilon))$ with $\epsilon \in \mathfrak{G}$, $\epsilon \prec 1$, $a \in \mathbb{R}$, $a > 0$. Let $\mathfrak{g} \in \mathfrak{G}$, $\mathfrak{g} \prec 1$. Then (i) $\epsilon(T) \sim \epsilon$; (ii) if $\mathfrak{g} \prec 1/(x\epsilon)$ then $\mathfrak{g}(T) \sim \mathfrak{g}$; (iii) if $\mathfrak{g} \sim -b/(x\epsilon)$, $b \in \mathbb{R}$ then $\mathfrak{g}(T) \sim e^{ab}\mathfrak{g}$; (iv) if $\mathfrak{g} \succ 1/(x\epsilon)$, then $\mathfrak{g}(T) \succ \mathfrak{g}$.*

Lemma 3.14. *Let $\mathfrak{m} \in \mathfrak{G}$ be a monomial. Let $\mathfrak{B} = \{\mathfrak{g} \in \mathfrak{G} : \mathfrak{g}^\dagger \prec \mathfrak{m}\}$ and $\tilde{\mathfrak{B}} = \{\mathfrak{g} \in \mathfrak{G} : \mathfrak{g}^\dagger \preccurlyeq \mathfrak{m}\}$. Then: (a) $\mathfrak{B}$ and $\tilde{\mathfrak{B}}$ are subgroups of $\mathfrak{G}$. (b) Let $\mathfrak{g} \in \mathfrak{G}$ be log-free and small. If $\mathfrak{g} \in \mathfrak{B}$, then $\text{supp}((x\mathfrak{g})') \subseteq \mathfrak{B}$. If $\mathfrak{g} \in \tilde{\mathfrak{B}}$, then $\text{supp}((x\mathfrak{g})') \subseteq \tilde{\mathfrak{B}}$. (c) Let $\mathfrak{g} = x^b e^L$, where L is purely large and log-free. If $\mathfrak{g} \in \mathfrak{B}$, then $\text{supp } L \subseteq \mathfrak{B}$. If $\mathfrak{g} \in \tilde{\mathfrak{B}}$, then $\text{supp } L \subseteq \tilde{\mathfrak{B}}$.*

Proof. (a) $(1/\mathfrak{g})^\dagger = -\mathfrak{g}^\dagger$, so if $\mathfrak{g} \in \mathfrak{B}$, then $1/\mathfrak{g} \in \mathfrak{B}$ and if $\mathfrak{g} \in \tilde{\mathfrak{B}}$, then $1/\mathfrak{g} \in \tilde{\mathfrak{B}}$. And $(\mathfrak{g}_1 \mathfrak{g}_2)^\dagger = \mathfrak{g}_1^\dagger + \mathfrak{g}_2^\dagger$, so if $\mathfrak{g}_1, \mathfrak{g}_2 \in \mathfrak{B}$, then $\mathfrak{g}_1 \mathfrak{g}_2 \in \mathfrak{B}$, and if $\mathfrak{g}_1, \mathfrak{g}_2 \in \tilde{\mathfrak{B}}$, then $\mathfrak{g}_1 \mathfrak{g}_2 \in \tilde{\mathfrak{B}}$.

(b) Apply Lemma 3.11(g).

(c) Let $\mathfrak{n} \in \text{supp } L$. Then by “height wins” we have $\mathfrak{n}^\dagger \preccurlyeq \mathfrak{g}^\dagger$. □

Lemma 3.15. G *Let $\mathfrak{A}$ be a subgrid. Then $\bigcup_{\mathfrak{a} \in \mathfrak{A}} \text{supp}(\mathfrak{a}')$ is also a subgrid. For any given $\mathfrak{g} \in \mathfrak{G}$, there are only finitely many $\mathfrak{a} \in \mathfrak{A}$ with $\mathfrak{g} \in \text{supp}(\mathfrak{a}')$.*

Proof. [11, Rem. 4.6]. (This is exactly what is needed for the proof that the derivative $(\sum_{\mathfrak{a} \in \mathfrak{A}} \mathfrak{a})'$ is defined.) □

Lemma 3.16. W *Let $\mathfrak{A}$ be well ordered. Then $\bigcup_{\mathfrak{a} \in \mathfrak{A}} \text{supp}(\mathfrak{a}')$ is also well ordered. For any given $\mathfrak{g} \in \mathfrak{G}$, there are only finitely many $\mathfrak{a} \in \mathfrak{A}$ with $\mathfrak{g} \in \text{supp}(\mathfrak{a}')$.*

Proof. [8, Lem. 3.2] or [10, Prop. 2.5]. □

Lemma 3.17. G *Let $\mathfrak{A}$ and $\mathfrak{B}$ be subgrids. Then $\mathfrak{A}\mathfrak{B}$ is a subgrid. If $\mathfrak{g} \in \mathfrak{A}\mathfrak{B}$, then there are only finitely many pairs $\mathfrak{a} \in \mathfrak{A}, \mathfrak{b} \in \mathfrak{B}$ with $\mathfrak{a}\mathfrak{b} = \mathfrak{g}$.*

Proof. [9, Prop. 3.35(d)] and [9, Prop. 3.27]. □

Lemma 3.18. W *Let $\mathfrak{A}$ and $\mathfrak{B}$ be well ordered. Then $\mathfrak{A}\mathfrak{B}$ is well ordered. If $\mathfrak{g} \in \mathfrak{A}\mathfrak{B}$, then there are only finitely many pairs $\mathfrak{a} \in \mathfrak{A}, \mathfrak{b} \in \mathfrak{B}$ with $\mathfrak{a}\mathfrak{b} = \mathfrak{g}$.*

Proof. [9, Prop. 3.27]. □

Lemma 3.19. G *Let $\mathfrak{B}$ be a subgrid. Let $\mathfrak{g} \in \mathfrak{G}$. There are only finitely many pairs $\mathfrak{g}_1, \mathfrak{g}_2 \in \mathfrak{B}$ such that $\mathfrak{g} \in \text{supp}((x\mathfrak{g}_1)'\mathfrak{g}_2)$.*

Proof. Since $\mathfrak{B}$ is a subgrid, also $\mathfrak{B}_1 := \{x\mathfrak{g} : \mathfrak{g} \in \mathfrak{B}\}$ is a subgrid. By Lemma 3.15, $\mathfrak{B}_2 := \bigcup_{\mathfrak{g} \in \mathfrak{B}} \text{supp}((x\mathfrak{g})')$ is a subgrid. Then $\mathfrak{B}_2\mathfrak{B} = \bigcup_{\mathfrak{g}_1, \mathfrak{g}_2 \in \mathfrak{B}} \text{supp}((x\mathfrak{g}_1)'\mathfrak{g}_2)$, and by Lemma 3.17, for any given $\mathfrak{g} \in \mathfrak{G}$, there are only finitely many pairs $\mathfrak{g}_1, \mathfrak{g}_2 \in \mathfrak{B}$ such that $\mathfrak{g} \in \text{supp}((x\mathfrak{g}_1)'\mathfrak{g}_2)$. □

Lemma 3.20. W *Let $\mathfrak{B}$ be well ordered. Let $\mathfrak{g} \in \mathfrak{G}$. There are only finitely many pairs $\mathfrak{g}_1, \mathfrak{g}_2 \in \mathfrak{B}$ such that $\mathfrak{g} \in \text{supp}((x\mathfrak{g}_1)'\mathfrak{g}_2)$.*

Proof. Since $\mathfrak{B}$ is well ordered, also $\mathfrak{B}_1 := \{x\mathfrak{g} : \mathfrak{g} \in \mathfrak{B}\}$ is well ordered. By Lemma 3.16, $\mathfrak{B}_2 := \bigcup_{\mathfrak{g} \in \mathfrak{B}} \text{supp}((x\mathfrak{g})')$ is well ordered, and each monomial in $\mathfrak{B}_2$ belongs to $\text{supp}((x\mathfrak{g})')$ for only finitely many $\mathfrak{g}$. Then $\mathfrak{B}_2\mathfrak{B} = \bigcup_{\mathfrak{g}_1, \mathfrak{g}_2 \in \mathfrak{B}} \text{supp}((x\mathfrak{g}_1)'\mathfrak{g}_2)$, and by Lemma 3.18, for any given $\mathfrak{g} \in \mathfrak{G}$, there are only finitely many pairs $\mathfrak{g}_2, \mathfrak{g}_1 \in \mathfrak{B}$ such that $\mathfrak{g} \in \text{supp}((x\mathfrak{g}_1)'\mathfrak{g}_2)$. □

Lemma 3.21. *Let $\mathfrak{e} \in \mathfrak{G}$, $\mathfrak{e} \prec 1$. Let*

$$\mathfrak{A} = \left\{ \mathfrak{g} \in \mathfrak{G} : \mathfrak{g} \preceq \mathfrak{e}, \mathfrak{g}^\dagger \prec \frac{1}{x\mathfrak{e}} \right\}, \quad \tilde{\mathfrak{A}} = \left\{ \mathfrak{g} \in \mathfrak{G} : \mathfrak{g} \preceq \mathfrak{e}, \mathfrak{g}^\dagger \preceq \frac{1}{x\mathfrak{e}} \right\}.$$

(a) *If $\mathfrak{g}_1, \mathfrak{g}_2 \in \mathfrak{A}$, then $\mathfrak{g}_1\mathfrak{g}_2 \in \mathfrak{A}$. If $\mathfrak{g}_1, \mathfrak{g}_2 \in \tilde{\mathfrak{A}}$, then $\mathfrak{g}_1\mathfrak{g}_2 \in \tilde{\mathfrak{A}}$.* (b) *If $\mathfrak{g}_1, \mathfrak{g}_2 \in \mathfrak{A}$, then $\text{supp}((x\mathfrak{g}_1)'\mathfrak{g}_2) \subseteq \mathfrak{A}$. If $\mathfrak{g}_1, \mathfrak{g}_2 \in \tilde{\mathfrak{A}}$, then $\text{supp}((x\mathfrak{g}_1)'\mathfrak{g}_2) \subseteq \tilde{\mathfrak{A}}$.* (c) *If $\mathfrak{g} \in \mathfrak{A}$, then $\text{supp}(x\mathfrak{e}\mathfrak{g}') \subseteq \mathfrak{A}$. If $\mathfrak{g} \in \tilde{\mathfrak{A}}$, then $\text{supp}(x\mathfrak{e}\mathfrak{g}') \subseteq \tilde{\mathfrak{A}}$.*

Proof. (a) If $\mathfrak{g}_1, \mathfrak{g}_2 \preceq \mathfrak{e}$, then $\mathfrak{g}_1\mathfrak{g}_2 \preceq \mathfrak{e}\mathfrak{e} \prec \mathfrak{e}$. Combine this with Lemma 3.14(a).

(b) If $\mathfrak{g}_1 \preceq \mathfrak{e}$, then $x\mathfrak{g}_1 \preceq x\mathfrak{e}$ and $(x\mathfrak{g}_1)' \preceq (x\mathfrak{e})'$. Now $\mathfrak{e} \prec 1$ so $x\mathfrak{e} \prec x$ and $(x\mathfrak{e})' \prec 1$. If $\mathfrak{g}_2 \preceq \mathfrak{e}$ also, then $(x\mathfrak{g}_1)'\mathfrak{g}_2 \preceq (x\mathfrak{e})'\mathfrak{e} \prec 1\mathfrak{e} = \mathfrak{e}$. Combine this with Lemma 3.14(b).

(c) is similar, noting that $\mathfrak{e}^\dagger \prec 1/(x\mathfrak{e})$ and $x^\dagger \prec 1/(x\mathfrak{e})$. $\square$

Lemma 3.22. $\boxed{\text{G}}$ *Let $\mathfrak{B} \subset \mathfrak{G}^{\text{small}}$ be a log-free subgrid. Write $\mathfrak{e} = \max \mathfrak{B}$ and assume $\mathfrak{B} \subseteq \left\{ \mathfrak{g} \in \mathfrak{G} : \mathfrak{g} \preceq \mathfrak{e}, \mathfrak{g}^\dagger \preceq 1/(x\mathfrak{e}) \right\}$. Let $\tilde{\mathfrak{B}}$ be the least subset of $\mathfrak{G}$ such that*

- (i) $\tilde{\mathfrak{B}} \supseteq \mathfrak{B}$,
- (ii) *if $\mathfrak{g}_1, \mathfrak{g}_2 \in \tilde{\mathfrak{B}}$, then $\mathfrak{g}_1\mathfrak{g}_2 \in \tilde{\mathfrak{B}}$,*
- (iii) *if $\mathfrak{g}_1, \mathfrak{g}_2 \in \tilde{\mathfrak{B}}$, then $\text{supp}((x\mathfrak{g}_1)'\mathfrak{g}_2) \subseteq \tilde{\mathfrak{B}}$.*

Then $\tilde{\mathfrak{B}}$ is a subgrid.

Proof. Let $\mathfrak{A}$ be the least subset of $\mathfrak{G}$ such that

- (i) $\mathfrak{A} \supseteq \mathfrak{B}$,
- (ii) *if $x^b e^L \in \mathfrak{A}$, then $\text{supp } L \subseteq \mathfrak{A}$.*

By [9, Prop. 2.21], $\mathfrak{A}$ is a subgrid. From Lemma 3.14(c) we have $\mathfrak{g}^\dagger \preceq 1/(x\mathfrak{e})$ for all $\mathfrak{g} \in \mathfrak{A}$. There is a ratio set $\boldsymbol{\mu} = \{\mu_1, \dots, \mu_n\}$, chosen from the group generated by $\mathfrak{A}$, so that $\mathfrak{A} \subseteq \mathfrak{J}^\mu$. Because they come from the group generated by $\mathfrak{A}$, we have $\mu_i^\dagger \preceq 1/(x\mathfrak{e})$ for $1 \leq i \leq n$ by Lemma 3.14(a). Remark that $x^\dagger \prec 1/(x\mathfrak{e})$ since $\mathfrak{e} \prec 1$. And $\mathfrak{e}^\dagger \prec 1/(x\mathfrak{e})$ was noted in Remark 3.4. So we may without harm add more generators to $\boldsymbol{\mu}$ and assume $\mathfrak{e}, x \in \mathfrak{J}^\mu$. This has been arranged so that if $\mathfrak{g} \in \mathfrak{J}^\mu$, then $\text{supp}(\mathfrak{g}') \subseteq \mathfrak{J}^\mu$. Now $x\mathfrak{e}\mu_i^\dagger \preceq 1$, so

$$\{\mathfrak{e}, x^{-1}\} \cup \bigcup_{i=1}^n \text{supp}(x\mathfrak{e}\mu_i^\dagger) \cup \frac{1}{\mathfrak{e}}\mathfrak{B}$$

is a finite union of subgrids, so it is itself a subgrid. All of its elements are $\preceq 1$, so [9, Prop. 3.52] there is a ratio set $\boldsymbol{\alpha}$ such that $\mathfrak{J}^{\boldsymbol{\alpha}, 0}$ contains that finite union. Again all elements of $\boldsymbol{\alpha}$ may be chosen from the group generated by $\mathfrak{J}^\mu$. So all $\mathfrak{a} \in \mathfrak{J}^{\boldsymbol{\alpha}, 0}$ still satisfy $\mathfrak{a}^\dagger \preceq 1/(x\mathfrak{e})$.

To complete the proof that $\tilde{\mathfrak{B}}$ is a subgrid, I will show that $\tilde{\mathfrak{B}} \subseteq \mathfrak{e}\mathfrak{J}^{\boldsymbol{\alpha}, 0}$. First, note that $\mathfrak{B} \subseteq \mathfrak{e}\mathfrak{J}^{\boldsymbol{\alpha}, 0}$. Next, if $\mathfrak{g}_1, \mathfrak{g}_2 \in \mathfrak{e}\mathfrak{J}^{\boldsymbol{\alpha}, 0}$, then $\mathfrak{g}_1\mathfrak{g}_2 \in \mathfrak{e}(\mathfrak{J}^{\boldsymbol{\alpha}, 0}\mathfrak{e}\mathfrak{J}^{\boldsymbol{\alpha}, 0}) \subseteq \mathfrak{e}\mathfrak{J}^{\boldsymbol{\alpha}, 0}$. Finally, suppose $\mathfrak{g}_1, \mathfrak{g}_2 \in \mathfrak{e}\mathfrak{J}^{\boldsymbol{\alpha}, 0}$. Because $\boldsymbol{\alpha}$ is from the group $\mathfrak{J}^\mu$, we may write $\mathfrak{g}_1 = \mu_1^{k_1} \cdots \mu_n^{k_n}$, and

$$\begin{aligned} \mathfrak{g}_1^\dagger &= k_1\mu_1^\dagger + \cdots + k_n\mu_n^\dagger, \\ x\mathfrak{e}\mathfrak{g}_1^\dagger &= k_1x\mathfrak{e}\mu_1^\dagger + \cdots + k_nx\mathfrak{e}\mu_n^\dagger, \end{aligned}$$

so that $\text{supp}(x\mathfrak{e}\mathfrak{g}_1^\dagger) \subseteq \mathfrak{J}^{\boldsymbol{\alpha}, 0}$. Also $\mathfrak{g}_1/\mathfrak{e} \in \mathfrak{J}^{\boldsymbol{\alpha}, 0}$ and $\mathfrak{g}_2/\mathfrak{e} \in \mathfrak{J}^{\boldsymbol{\alpha}, 0}$. Therefore

$$\text{supp}(x\mathfrak{g}_1'\mathfrak{g}_2) = \mathfrak{e} \left(\frac{\mathfrak{g}_1}{\mathfrak{e}} \right) \left(\frac{\mathfrak{g}_2}{\mathfrak{e}} \right) \text{supp}(x\mathfrak{e}\mathfrak{g}_1^\dagger) \subseteq \mathfrak{e}\mathfrak{J}^{\boldsymbol{\alpha}, 0}.$$

And $(xg_1)'g_2 = g_1g_2 + xg_1'g_2$, so $\text{supp}((xg_1)'g_2) \subseteq \mathfrak{e}\mathfrak{J}^{\alpha,0}$. By the definition of $\tilde{\mathfrak{B}}$ we have $\tilde{\mathfrak{B}} \subseteq \mathfrak{e}\mathfrak{J}^{\alpha,0}$, and it is therefore a subgrid. $\square$

Lemma 3.23. W *Let $\mathfrak{B} \subset \mathfrak{G}^{\text{small}}$ be log-free and well ordered. Write $\mathfrak{e} = \max \mathfrak{B}$ and assume $\mathfrak{B} \subseteq \{g \in \mathfrak{G} : g \preccurlyeq \mathfrak{e}, g^\dagger \preccurlyeq 1/(x\mathfrak{e})\}$. Let $\tilde{\mathfrak{B}}$ be the least subset of $\mathfrak{G}$ such that*

- (i) $\tilde{\mathfrak{B}} \supseteq \mathfrak{B}$,
- (ii) if $g_1, g_2 \in \tilde{\mathfrak{B}}$, then $g_1g_2 \in \tilde{\mathfrak{B}}$,
- (iii) if $g_1, g_2 \in \tilde{\mathfrak{B}}$, then $\text{supp}((xg_1)'g_2) \subseteq \tilde{\mathfrak{B}}$.

Then $\tilde{\mathfrak{B}}$ is well ordered.

Proof. Let $\mathfrak{B}_1$ be the least set such that $\mathfrak{B}_1 \supseteq \mathfrak{B} \cup \{\mathfrak{e}^2\}$ and if $g \in \mathfrak{B}_1$, then $\text{supp}(xeg') \subseteq \mathfrak{B}_1$. Then $\mathfrak{B}_1$ is well ordered by [10, Prop. 2.10]. For all $g \in \mathfrak{B}_1$ we have $g \preccurlyeq \mathfrak{e}$ and $g^\dagger \preccurlyeq 1/(x\mathfrak{e})$ by Lemma 3.21(c). Still $\mathfrak{e} = \max \mathfrak{B}_1$, $\mathfrak{e}^2 \in \mathfrak{B}_1$, and $\text{supp}(x\mathfrak{e}\mathfrak{e}') \subseteq \mathfrak{B}_1$.

Let $\mathfrak{B}_2 = \mathfrak{e}^{-1}\mathfrak{B}_1$. Then $\mathfrak{B}_2$ is well ordered, $\mathfrak{B}_2 \supseteq \mathfrak{e}^{-1}\mathfrak{B}$, $1 = \max \mathfrak{B}_2$, $\mathfrak{e} \in \mathfrak{B}_2$, $\text{supp}(x\mathfrak{e}') \subseteq \mathfrak{B}_2$. If $m \in \mathfrak{B}_2$, then $\text{supp}(x(\mathfrak{e}m)') \subseteq \mathfrak{B}_2$.

Let $\mathfrak{B}_3$ be the semigroup generated by $\mathfrak{B}_2$. Then $\mathfrak{B}_3$ is well ordered, $\mathfrak{B}_3 \supseteq \mathfrak{e}^{-1}\mathfrak{B}$, $1 = \max \mathfrak{B}_3$, $\mathfrak{e} \in \mathfrak{B}_3$, $\text{supp}(x\mathfrak{e}') \subseteq \mathfrak{B}_3$. From the identity

$$x(\mathfrak{e}m_1m_2)' = x(\mathfrak{e}m_1)' \cdot m_2 + m_1 \cdot x(\mathfrak{e}m_2)' - x\mathfrak{e}'m_1m_2$$

we conclude: if $m \in \mathfrak{B}_3$, then $\text{supp}(x(\mathfrak{e}m)') \subseteq \mathfrak{B}_3$.

Finally, let $\mathfrak{B}_4 = \mathfrak{e}\mathfrak{B}_3$. Then $\mathfrak{B}_4$ is well ordered, $\mathfrak{B}_4 \supseteq \mathfrak{B}$, $\mathfrak{e} = \max \mathfrak{B}_4$. Let $g_1, g_2 \in \mathfrak{B}_4$. Then $g_1/\mathfrak{e}, g_2/\mathfrak{e} \in \mathfrak{B}_3$, so $(g_1/\mathfrak{e}) \cdot (g_2/\mathfrak{e}) \in \mathfrak{B}_3$ and $g_1g_2/\mathfrak{e}^2 \in \mathfrak{B}_3$. Now $\mathfrak{e} \in \mathfrak{B}_3$ so $g_1g_2/\mathfrak{e} \in \mathfrak{B}_3$ and therefore $g_1g_2 \in \mathfrak{B}_4$. Again let $g_1, g_2 \in \mathfrak{B}_4$. Then $g_1/\mathfrak{e}, g_2/\mathfrak{e} \in \mathfrak{B}_3$. So $\text{supp}(xg_1') \subseteq \mathfrak{B}_3$. Thus $\text{supp}(xg_1')g_2/\mathfrak{e} \subseteq \mathfrak{B}_3$ so $\text{supp}(xg_1')g_2 \subseteq \mathfrak{B}_4$. And $(xg_1)'g_2 = g_1g_2 + xg_1'g_2$, so we conclude $\text{supp}((xg_1)'g_2) \subseteq \mathfrak{B}_4$.

This shows $\mathfrak{B}_4 \supseteq \tilde{\mathfrak{B}}$, and therefore that $\tilde{\mathfrak{B}}$ is well ordered. $\square$

4 Abel's Equation

Let $T \in \mathcal{P}$, $T > x$. **Abel's Equation** for T is $V(T(x)) = V(x) + 1$. If large positive V exists satisfying this, then a real iteration group Φ may be obtained as $\Phi(s, x) = V^{[-1]} \circ (x + s) \circ V$. (In general such Φ will not have common support.) If $T \in \mathcal{P}$, $T < x$, then Abel's Equation for T is $V(T(x)) = V(x) - 1$, and then we may similarly write $\Phi(s, x) = V^{[-1]} \circ (x - s) \circ V$.

We now do this in reverse: Let $\Phi(s, x)$ be of the form constructed as in Theorem 3.8, that is, coefficients defined recursively by (15) and (17). Then we can use Φ to get V for Abel's Equation.

Theorem 4.1. (Moderate Abel) *Let $T \sim x$, $T > x$ be moderate, and let $\Phi(s, x) \in \mathcal{P}$ for all $s \in \mathbb{R}$ be the real iteration group for T constructed in Theorem 3.8. Then*

$$V(x) := \int \frac{dx}{\Phi_1(0, x)}$$

is large and positive and satisfies Abel's Equation $V(T(x)) = V(x) + 1$.

Proof. Now $V' \sim 1/(ax\epsilon) \succ 1/x$, so $V \succ \log x$ is large. And $V' > 0$ so $V > 0$. (A large negative transseries has negative derivative.)

From $\Phi(s+t, x) = \Phi(s, \Phi(t, x))$ take $\partial/\partial s$ then substitute $t = 1, s = 0$ to get $\Phi_1(1, x) = \Phi_1(0, T)$. As constructed, $\Phi_1(s, x) = \Phi_2(s, x)\Phi_1(0, x)$. So $\Phi_2(1, x)\Phi_1(0, x) = \Phi_1(1, x) = \Phi_1(0, T)$. Now from $\Phi(1, x) = T$ we have $\Phi_2(1, x) = T'$. So

$$\frac{T'}{\Phi_1(0, T)} = \frac{1}{\Phi_1(0, x)},$$

or $V'(T) \cdot T' = V'(x)$ so $V(T) = V + c$ for some $c \in \mathbb{R}$. Now

$$V(T) - V(x) = \int_x^T V' = \int_x^T \frac{1}{\Phi_1(0, x)} \sim \int_x^T \frac{1}{ax\epsilon}.$$

By [10, Prop. 4.24], this integral is between

$$\frac{T-x}{ax\epsilon} \sim \frac{ax\epsilon}{ax\epsilon} = 1 \quad \text{and} \quad \frac{T-x}{aT\epsilon(T)} \sim \frac{ax\epsilon}{ax\epsilon} = 1.$$

We used $\epsilon(T) \sim \epsilon$ from Lemma 3.12(i). So we have $V(T) - V \sim 1$ and thus $c = 1$. $\square$

Now we will consider the deep case. For $T = x + 1 + A$, consider Abel's Equation $V \circ T = V + 1$. A formal solution is

$$V = x + A + A \circ T + A \circ T^{[2]} + A \circ T^{[3]} + \dots. \quad (18)$$

But if T is not purely deep, then $A \circ T \asymp A$ (Lemma 3.13), so series (18) does not converge. We will use the moderate version already proved (Theorem 4.1) to reduce the general case to one where $A^\dagger \succ 1$ (Proposition 4.2) so that $A \circ T \prec A$ and the series does converge (Proposition 4.3). But in general it cannot be grid-based (Example 4.8), so the final step works only for the well-based version of $\mathbb{T}$.

Proposition 4.2. *Let $T \sim x$, $T > x$. There exists large positive V such that $V \circ T \circ V^{[-1]} = x + 1 + B$ and $\mathfrak{g}^\dagger \succ 1$ for all $\mathfrak{g} \in \text{supp } B$; that is, $x + 1 + B$ is purely deep. Let $0 < S < x$, $S \succ 1$. There exists $V \in \mathcal{P}$ such that $V \circ S \circ V^{[-1]} = x - 1 + C$ and $\mathfrak{g}^\dagger \succ 1$ for all $\mathfrak{g} \in \text{supp } C$.*

Proof. Write $T = x(1 + a\epsilon + A_1 + A_2)$, where

- (i) $\mathfrak{g} \prec \epsilon, \mathfrak{g}^\dagger \preceq 1/(x\epsilon)$ for all $\mathfrak{g} \in \text{supp } A_1$,
- (ii) $\mathfrak{g} \prec 1, \mathfrak{g}^\dagger \succ 1/(x\epsilon)$ for all $\mathfrak{g} \in \text{supp } A_2$.

So $T_1 = x(1 + a\epsilon + A_1)$ is the moderate part of T (including the shallow part), and $T - T_1 = xA_2$ is the deep part of T . By Theorem 4.1, there is large positive $V \succ \log x$ so that $V \circ T_1 = V + 1$ and $V' \sim 1/(ax\epsilon)$. So compute

$$V \circ T - V \circ T_1 = \int_{T_1}^T V' \succ \int_{T_1}^T \frac{1}{x}.$$

Now $T - T_1 = A_2$, so this integral is between $A_2/T_1 \sim A_2/x$ and $A_2/T \sim A_2/x$. So $B_1 := V \circ T - V \circ T_1 \prec A_2$ and

$$V \circ T = V \circ T_1 + B_1 = V + 1 + B_1$$

with $B_1^\dagger \succ 1/(x\epsilon)$. So write $B = B_1 \circ V^{[-1]}$ to get $V \circ T \circ V^{[-1]} = x + 1 + B$ and $B_1^\dagger = (B \circ V)^\dagger = (B^\dagger \circ V) \cdot V'$. But $V' \asymp 1/(x\epsilon)$ and $B_1^\dagger \succ 1/(x\epsilon)$, so $B^\dagger \succ 1$.

Now let $0 < S < x$, $S \succ 1$. Then define $T := S^{[-1]}$ to get $T > x$, $T \succ 1$. So as we have just seen, there is V with $V \circ T \circ V^{[-1]} = x + 1 + B$. Take the inverse to get

$$V \circ S \circ V^{[-1]} = (V \circ T \circ V^{[-1]})^{[-1]} = (x + 1 + B)^{[-1]}.$$

So if $(x+1+B)^{[-1]} = (x-1+C)$, we must show $C^\dagger \succ 1$. Now $(x+1+B) \circ (x-1+C) = x$, so $x-1+C+1+B \circ (x-1+C) = x$ and therefore $C = -B \circ (x-1+C)$, so

$$C^\dagger = (B^\dagger \circ (x-1+C)) \cdot (x-1+C)' \succ 1 \cdot 1 = 1,$$

as required. $\square$

The following proof is only for the well-based version of $\mathbb{T}$. In Example 4.8, below, we see it fails in general for the grid-based version of $\mathbb{T}$.

Proposition 4.3. $\boxed{\text{W}}$ (Purely Deep Abel) (a) Let $T = x + 1 + A$, $A \in \mathbb{T}$, $A \prec 1$, $A^\dagger \succ 1$. There is $V = x + B$, $B \in \mathbb{T}$, $B \prec 1$, $B^\dagger \succ 1$, such that $V \circ T = V + 1$. (b) Let $T = x - 1 + A$, $A \in \mathbb{T}$, $A \prec 1$, $A^\dagger \succ 1$. There is $V = x + B$, $B \in \mathbb{T}$, $B \prec 1$, $B^\dagger \succ 1$ such that $V \circ T = V - 1$.

Proof. (a) There exist $N, M \in \mathbb{N}$ so that $\text{supp } A \subseteq \mathfrak{G}_{N,M}$. Increase N if necessary so that $N \geq M$ and $x \in \mathfrak{G}_{N,M}$. Now $\epsilon = x^{-1}$, so the deep monomials are $\mathfrak{D} = \{ \mathfrak{g} \in \mathfrak{G}_{N,M}^{\text{small}} : \mathfrak{g}^\dagger \succ 1 \}$. I claim: if $B \in \mathbb{T}$, $\text{supp } B \subseteq \mathfrak{D}$, then $B(T) \prec B$ and $\text{supp } B(T) \subseteq \mathfrak{D}$. Indeed, all $\mathfrak{g} \in \text{supp } B$ satisfy $\mathfrak{g}(T) \prec \mathfrak{g}$ by Lemma 3.12(iv), and we may sum to conclude $B(T) \prec B$. Therefore $B(T)^\dagger \succ B^\dagger \succ 1$. And $\text{supp } B(T) \subseteq \mathfrak{G}_{N,M}$ by [9, Prop. 3.111].

Let $\mathcal{A} = \{ x + B \in \mathbb{T} : \text{supp } B \subseteq \mathfrak{D} \}$. Define Ψ by $\Psi(Y) := Y \circ T - 1$. We want to apply a fixed point argument to Ψ . First we must show that Ψ maps $\mathcal{A}$ into $\mathcal{A}$. Let $x + B \in \mathcal{A}$. So $\Psi(x + B) = T + B \circ T - 1 = x + 1 + A + B(T) - 1$. But $\text{supp } A \subseteq \mathfrak{D}$, $\text{supp } B \subseteq \mathfrak{D}$ so $\text{supp } B(T) \subseteq \mathfrak{D}$, and thus $x + A + B(T) \in \mathcal{A}$.

Suppose $x + B_1, x + B_2 \in \mathcal{A}$. Then $\text{supp}(B_1 - B_2) \subseteq \mathfrak{D}$ and

$$\begin{aligned} \Psi(x + B_1) - \Psi(x + B_2) &= (T + B_1 \circ T - 1) - (T + B_2 \circ T - 1) \\ &= (B_1 - B_2) \circ T \prec B_1 - B_2. \end{aligned}$$

So Ψ is contractive.

Now we are ready to apply the well based contraction theorem [14, Thm 4.7]. In our case where $\mathfrak{G}$ is totally ordered, the dotted ordering $\prec$ of [14] coincides with the usual ordering $\prec$. There is $V \in \mathcal{A}$ such that $\Psi(V) = V$. This is what was required.

Part (b) is proved from part (a) as before: Begin with $T = x - 1 + A$ purely deep, then $T^{[-1]} = x + 1 + A_1$ also purely deep, from part (a) get $V = x + B$ with $V \circ T^{[-1]} = V + 1$, so compose with T on the right to get $V = V \circ T + 1$ as desired. $\square$

Because they depend on Proposition 4.3, the following two results are also valid only for the well-based version of $\mathbb{T}$.

Theorem 4.4. $\boxed{\text{W}}$ (General Abel) Let $T \in \mathcal{P}$ with $\text{expo } T = 0$. Then there is $V \in \mathcal{P}$ such that: (i) If $T > x$, then $V \circ T \circ V^{[-1]} = x + 1$; (ii) If $T < x$, then $V \circ T \circ V^{[-1]} = x - 1$.

Proof. (i) First, there is V_1 so that $T_1 := V_1 \circ T \circ V_1^{[-1]} \sim x$. By Proposition 4.2, there is V_2 so that $T_2 := V_2 \circ T_1 \circ V_2^{[-1]} = x + 1 + B$ with $B^\dagger \prec 1$. By Proposition 4.3 there is V_3 so that $V_3 \circ T_2 \circ V_3^{[-1]} = x + 1$. Define $V = V_3 \circ V_2 \circ V_1$ to get $V \circ T \circ V^{[-1]} = x + 1$.
(ii) is similar. $\square$

Corollary 4.5. $\boxed{\text{W}}$ *Let $T \in \mathcal{P}$ with $\text{expo } T = 0$. Then there exists real iteration group $\Phi(s, x)$ for T .*

Proof. In case $T > x$, let V be as in Theorem 4.4(i), then take $\Phi(s, x) = V^{[-1]} \circ (x + s) \circ V$. In case $T < x$, let V be as in Theorem 4.4(ii), then take $\Phi(s, x) = V^{[-1]} \circ (x - s) \circ V$. $\square$

Question 4.6. The proof as given here depends on the existence of inverses in $\mathcal{P}$. Is it possible to demonstrate first the solution to Abel's Equation without assuming the existence of inverses, then use that to construct inverses?

Example 4.7. Take the example $T = x + 1 + xe^{-x^2}$ of Proposition 2.6. Carrying out the iteration of Proposition 4.3, we get V satisfying $V(T(x)) = V(x) + 1$ which looks like:

$$\begin{aligned}
V = & x \\
& + e^{-x^2} \left(x + e^{-2x}(x+1)e^{-1} + e^{-4x}(x+2)e^{-4} + e^{-6x}(x+3)e^{-9x} + \dots \right) \\
& + e^{-2x^2} \left(e^{-2x}(-x - 4x^2 - 2x^3)e^{-1} + e^{-4x}(x - 4x^2 - 2x^3)e^{-4} \right. \\
& \quad \left. + e^{-6x}[(7x - 4x^2 - 2x^3)e^{-9} + (-7 - 15x - 10x^2 - 2x^3)e^{-5}] + \dots \right) \\
& + e^{-3x^2} \left(e^{-2x}(-x^2 + 3x^3 + 6x^4 + 2x^5)e^{-1} + e^{-4x}(-2x^2 - 3x^3 + 4x^4 + 2x^5)e^{-4} \right. \\
& \quad \left. + e^{-6x}[(+5x^2 - 13x^3 + 2x^4 + 2x^5)e^{-9} \right. \\
& \quad \left. + (-x + 38x^2 + 74x^3 + 44x^4 + 8x^5)e^{-5}] + \dots \right) \\
& + e^{-4x^2} \left(e^{-2x} \left(\frac{5}{3}x^3 + \frac{8}{3}x^4 - 4x^5 - \frac{16}{3}x^6 - \frac{4}{3}x^7 \right) e^{-1} \right. \\
& \quad \left. + e^{-4x} \left(-x^3 + 4x^4 + 4x^5 - \frac{8}{3}x^6 - \frac{4}{3}x^7 \right) e^{-4} + \dots \right) \\
& + \dots
\end{aligned}$$

The support is a subgrid of order type ω^2 .

Of course, once we have V we can compute the real iteration group $\Phi(s, x) = V^{[-1]} \circ (x + s) \circ V$. For s negative we get

$$\Phi(s, x) = x + s - xe^{-(x+s)^2} + \dots,$$

so they are not contained in a common grid (or well ordered set), as noted before. But since V is grid-based, all of the fractional iterates $T^{[s]}$ are also grid-based.

The Non-Grid Situation

Example 4.8. Here is an example where Abel's Equation has no grid-based solution.

$$T = x + 1 + e^{-e^{x^2}}.$$

The support for V where $V \circ T = V + 1$ deserves careful examination. We will use these notations:

$$\begin{aligned} \mathfrak{m}_1 &= x^{-1}, & \mathfrak{m}_1 &\prec 1, & \mathfrak{m}_1 &\in \mathfrak{G}_0, \\ \mathfrak{m}_2 &= e^{-x}, & \mathfrak{m}_2 &\prec \mathfrak{m}_1, & \mathfrak{m}_2 &\in \mathfrak{G}_1, \\ \mathfrak{m}_3 &= e^{-x^2}, & \mathfrak{m}_3 &\prec \mathfrak{m}_2, & \mathfrak{m}_3 &\in \mathfrak{G}_1, \\ \boldsymbol{\mu} &= \{\mathfrak{m}_1, \mathfrak{m}_2, \mathfrak{m}_3\} \subset \mathfrak{G}_1, \\ L_k &= e^{(x+k)^2} = e^{k^2} \mathfrak{m}_2^{-2k} \mathfrak{m}_3^{-1}, & k &= 0, 1, 2, \dots, & \text{supp } L_k &\subset \mathfrak{J}^\mu \subset \mathfrak{G}_1, \\ \mathfrak{a}_k &= e^{-L_k}, & k &= 0, 1, 2, \dots, & \mathfrak{a}_k &\in \mathfrak{G}_2, \\ \mathfrak{b}_k &= x \mathfrak{m}_2^{-2k} \mathfrak{m}_3^{-1} \mathfrak{a}_k, & \boldsymbol{\alpha} &= \boldsymbol{\mu} \cup \{\mathfrak{b}_k : k = 0, 1, 2, \dots\} \subset \mathfrak{G}_2. \end{aligned}$$

Now $\boldsymbol{\alpha}$ is infinite, so it is not a ratio set in the usual sense. However, writing $\mathfrak{g}_1 \succ \mathfrak{g}_2$ iff $\mathfrak{g}_1^k \succ \mathfrak{g}_2$ for all $k \in \mathbb{N}$, we have

$$\mathfrak{m}_1 \succ \mathfrak{m}_2 \succ \mathfrak{m}_3 \succ \mathfrak{b}_0 \succ \mathfrak{b}_1 \succ \mathfrak{b}_2 \succ \dots.$$

The semigroup generated by $\boldsymbol{\alpha}$ is contained in $\mathfrak{G}_2$, is well ordered, and has order type ω^ω . Probably the solution V of Abel's Equation also has support of order type ω^ω , but to prove it we would have to verify that many terms are not eliminated by cancellation.

Computations follow. When I write $\mathfrak{o}$ and $\mathfrak{O}$, the omitted terms all belong to $\mathfrak{G}_2$.

$$\begin{aligned} T &= x + 1 + \mathfrak{a}_0, \\ T^2 &= x^2 + 2x + 1 + 2x\mathfrak{a}_0 + \mathfrak{O}(\mathfrak{a}_0), \\ \mathfrak{m}_2^{-1} \circ T &= e^T = e^{x+1+\mathfrak{a}_0} = e\mathfrak{m}_2^{-1}e^{\mathfrak{a}_0} = e\mathfrak{m}_2^{-1}(1 + \mathfrak{a}_0 + \mathfrak{o}(\mathfrak{a}_0)) \\ &= e\mathfrak{m}_2^{-1} + e\mathfrak{m}_2^{-1}\mathfrak{a}_0 + \mathfrak{o}(\mathfrak{m}_2^{-1}\mathfrak{a}_0), \\ \mathfrak{m}_2^{-2k} \circ T &= e^{2k} \mathfrak{m}_2^{-2k} + 2ke^{2k} \mathfrak{m}_2^{-2k} \mathfrak{a}_0 + \mathfrak{o}(\mathfrak{m}_2^{-2k} \mathfrak{a}_0), \\ \mathfrak{m}_3^{-1} \circ T &= e^{T^2} = e^{x^2+2x+1+2x\mathfrak{a}_0+\mathfrak{O}(\mathfrak{a}_0)} = e\mathfrak{m}_2^{-2}\mathfrak{m}_3^{-1}e^{2x\mathfrak{a}_0+\mathfrak{O}(\mathfrak{a}_0)} \\ &= e\mathfrak{m}_2^{-2}\mathfrak{m}_3^{-1}(1 + 2x\mathfrak{a}_0 + \mathfrak{O}(\mathfrak{a}_0)) \\ &= e\mathfrak{m}_2^{-2}\mathfrak{m}_3^{-1} + 2ex\mathfrak{m}_2^{-2}\mathfrak{m}_3^{-1}\mathfrak{a}_0 + \mathfrak{O}(\mathfrak{m}_2^{-2}\mathfrak{m}_3^{-1}\mathfrak{a}_0), \\ L_k \circ T &= e^{k^2+2k+1} \mathfrak{m}_2^{-2k-2} \mathfrak{m}_3^{-1} + 2e^{k^2+2k+1} x \mathfrak{m}_2^{-2k-2} \mathfrak{m}_3^{-1} \mathfrak{a}_0 + \mathfrak{O}(\mathfrak{m}_2^{-2k-2} \mathfrak{m}_3^{-1} \mathfrak{a}_0) \\ \mathfrak{a}_k \circ T &= e^{-L_{k+1}} e^{-2e^{(k+1)^2} x \mathfrak{m}_2^{-2k-2} \mathfrak{m}_3^{-1} \mathfrak{a}_0 + \mathfrak{O}(\mathfrak{m}_2^{-2k-2} \mathfrak{m}_3^{-1} \mathfrak{a}_0)} \\ &= \mathfrak{a}_{k+1} (1 - 2e^{(k+1)^2} x \mathfrak{m}_2^{-2k-2} \mathfrak{m}_3^{-1} \mathfrak{a}_0 + \mathfrak{O}(\mathfrak{m}_2^{-2k-2} \mathfrak{m}_3^{-1} \mathfrak{a}_0)) \\ &= \mathfrak{a}_{k+1} - 2e^{(k+1)^2} x \mathfrak{m}_2^{-2k-2} \mathfrak{m}_3^{-1} \mathfrak{a}_0 \mathfrak{a}_{k+1} + \mathfrak{O}(\mathfrak{m}_2^{-2k-2} \mathfrak{m}_3^{-1} \mathfrak{a}_0 \mathfrak{a}_{k+1}) \\ &= \mathfrak{a}_{k+1} - 2e^{(k+1)^2} \mathfrak{a}_0 \mathfrak{b}_{k+1} + \mathfrak{o}(\mathfrak{a}_0 \mathfrak{b}_{k+1}) \\ \mathfrak{b}_k \circ T &= (x \mathfrak{m}_2^{-2k} \mathfrak{m}_3^{-1} \mathfrak{a}_k) \circ T = \mathfrak{b}_{k+1} + \mathfrak{o}(\mathfrak{b}_{k+1}). \end{aligned}$$

The solution V of Abel's Equation $V \circ T = V + 1$ is

$$V = x + 1 + \mathfrak{a}_0 + \mathfrak{a}_0 \circ T + \mathfrak{a}_0 \circ T^{[2]} + \mathfrak{a}_0 \circ T^{[3]} + \dots.$$

Without considering cancellation, we would expect that its support still has order type ω^ω . Even without trying to account for cancellation, we know that $\text{supp } V$ contains $\{\mathfrak{a}_0, \mathfrak{a}_1, \mathfrak{a}_2, \dots\}$. The logarithms L_k are linearly independent, so the group generated by $\{\mathfrak{a}_k : k \in \mathbb{N}\}$ is not finitely generated, and thus $\text{supp } V$ is not a subgrid. $\square$

More computation in this example yields: $V^{[-1]} = x - 1 - \mathfrak{a}_{-1} + \dots$ and

$$T^{[1/2]} = V^{[-1]} \circ \left(x + \frac{1}{2}\right) \circ V = x + \frac{1}{2} + \mathfrak{a}_0 - \mathfrak{a}_{1/2} + \dots$$

not grid-based. We used notation $\mathfrak{a}_k = \exp(-\exp((x+k)^2))$ for $k = -1$ and $1/2$.

Consider the proof of Proposition 4.3. How much can be done in the grid-based version? Assume grid-based $T = x + 1 + A$, $A \in \mathbb{T}$, $A \prec 1$, $A^\dagger \succ 1$. Write $\mathfrak{m} = \text{mag } A$. Consider a ratio set μ such that $\text{supp } A \subseteq \mathfrak{m}\mathfrak{J}^{\mu,0}$. There is [11, Prop. 5.6] a “ T -composition addendum” α for μ such that:

- (i) if $\mathfrak{a} \in \mathfrak{J}^{\mu,0}$, then $\text{supp}(\mathfrak{a} \circ T) \subseteq \mathfrak{J}^{\alpha,0}$;
- (ii) if $\mathfrak{a} \prec^\mu \mathfrak{b}$, then $\mathfrak{a} \circ T \prec^\alpha \mathfrak{b} \circ T$.

But this is not enough to carry out the contraction argument. We need a “hereditary T -composition addendum” $\alpha \supseteq \mu$ such that:

- (i) if $\mathfrak{a} \in \mathfrak{J}^{\alpha,0}$, then $\text{supp}(\mathfrak{a} \circ T) \subseteq \mathfrak{J}^{\alpha,0}$;
- (ii) if $\mathfrak{a} \prec^\alpha \mathfrak{b}$, then $\mathfrak{a} \circ T \prec^\alpha \mathfrak{b} \circ T$.

For some deep T there is such an addendum, but not for others. If there is, then a grid-based version of the contraction argument of Proposition 4.3 works. Or (for purely deep T) we can write

$$V = x + 1 + A + A \circ T + A \circ T^{[2]} + A \circ T^{[3]} + \dots$$

with $A \succ^\alpha A \circ T \succ^\alpha A \circ T^{[2]} \succ^\alpha \dots$ to insure grid-based convergence in the asymptotic topology.

For example, if $\mu = \{x^{-1}, e^{-x}, e^{-x^2}\}$ and $\text{supp } A \subseteq \mathfrak{J}^{\mu,0}$, then $\alpha = \{x^{-1}, e^{-x}, x^2 e^{-x^2}\}$ is a hereditary T -composition addendum. This insures that the iteration used in Example 4.7 provides a grid-based solution V .

5 Uniqueness

In what sense is $T^{[s]}$ unique? This question is related to the question of commutativity for composition.

Proposition 5.1. *Let $V \in \mathcal{P}$. If $V(x+1) = V+1$, then there is $c \in \mathbb{R}$ with $V = x+c$.*

Proof. By [10, Prop. 4.23], $1 = (V(x+1) - V(x))/1 = V' \circ S$ for some $S \in \mathcal{P}$. Compose on the right with $S^{[-1]}$ to get $1 = V'$. So $V = x+c$ as required. $\square$

Corollary 5.2. *Let $T \in \mathcal{P}$, $T > x$. The solution $V \in \mathcal{P}$ of $V \circ T = V+1$ is unique up to a constant addend.*

Proof. Suppose $V \circ T = V+1$ and $U \circ T = U+1$. Then $V^{[-1]} \circ (x+1) \circ V = U^{[-1]} \circ (x+1) \circ U$ and $(U \circ V^{[-1]}) \circ (x+1) = (x+1) \circ (U \circ V^{[-1]})$. By Proposition 5.1 there is $c \in \mathbb{T}$ with $U \circ V^{[-1]} = x+c$ so that $U = V+c$. $\square$

Notation 5.3. Let $T \in \mathcal{P}$, $s \in \mathbb{R}$. If $T > x$, define $T^{[s]} = V^{[-1]} \circ (x + s) \circ V$, where V is a solution of Abel's Equation $V(T) = V + 1$. If $T < x$, define $T^{[s]} = V^{[-1]} \circ (x - s) \circ V$, where V is a solution of Abel's Equation $V(T) = V - 1$. The transseries $T^{[s]}$ is independent of the choice of solution V .

Note: Even if T is grid-based, it could happen that $T^{[s]}$ is not.

Proposition 5.4. *Let $A, B \in \mathcal{P}$, $B \neq x$. If $A \circ B = B \circ A$, then there is $s \in \mathbb{R}$ with $B^{[s]} = A$.*

Proof. We do the case $B > x$; the case $B < x$ is similar. Let $V \in \mathcal{P}$ solve Abel's Equation for B , so that $B^{[s]} = V^{[-1]} \circ (x + s) \circ V$ for $s \in \mathbb{R}$. Then $V^{[-1]} \circ (x + 1) \circ V \circ A = A \circ V^{[-1]} \circ (x + 1) \circ V$. Compose with V on the left and $V^{[-1]}$ on the right to get $(x + 1) \circ (V \circ A \circ V^{[-1]}) = (V \circ A \circ V^{[-1]}) \circ (x + 1)$. By Proposition 5.1, there is $s \in \mathbb{R}$ with $V \circ A \circ V^{[-1]} = x + s$. So $A = V^{[-1]} \circ (x + s) \circ V = B^{[s]}$.

If A, B are grid-based, perhaps $B^{[s]}$ is in general not grid-based. But since we conclude $B^{[s]} = A$, then at least for this particular s it happens to be grid-based. $\square$

Example 5.5. Let $\theta: \mathbb{R} \rightarrow \mathbb{R}$ satisfy $\theta(1) = 1$ and $\theta(s+t) = \theta(s) + \theta(t)$ for all s, t . By the Axiom of Choice, there is such a map θ other than the identity function $\theta(s) = s$. (This strange θ is everywhere discontinuous, non-measurable, unbounded on every interval.) Let $T \in \mathcal{P}$. Then $\Phi(s, x) = T^{[\theta(s)]}$ is a real iteration group for T .

Here is a way to rule out such strange cases.

Proposition 5.6. *Let $T \in \mathcal{P}$, $T > x$, and let $\Phi(s, x)$ be a real iteration group for T . Assume $\Phi(s, x) > x$ for all $s > 0$. Then $\Phi(s, x) = T^{[s]}$ as in 5.3.*

Proof. Since $\Phi(s, x) > x$ for $s > 0$, we may deduce that $s_1 < s_2$ implies $\Phi(s_1, x) < \Phi(s_2, x)$. Also $\Phi(1, x) = T$, so we may deduce $\Phi(s, x) = T^{[s]}$ for all rational s . Fix an irrational s . Since $\Phi(s, x) \circ \Phi(1, x) = \Phi(s+1, x)$, we know that $\Phi(s, x)$ commutes with T , so by Proposition 5.4, $\Phi(s, x) = T^{[t]}$ for some t . But the only t satisfying $T^{[s_1]} < T^{[t]} < T^{[s_2]}$ for all rationals s_1, s_2 with $s_1 < s < s_2$ is $t = s$ itself. $\square$

Similarly: Let $T \in \mathcal{P}$, $T < x$, and let $\Phi(s, x)$ be a real iteration group for T . Assume $\Phi(s, x) < x$ for all $s > 0$. Then $\Phi(s, x) = T^{[s]}$ as in 5.3.

6 Julia Example

As an example we will consider fractional iterates for the function $M(x) = x^2 + c$ near $x = +\infty$. Of course, integer iterates of this function are used for construction of Julia sets or the Mandelbrot set. For the theory of real transseries to be applicable, we must restrict to real values c . But once we have nice formulas, they can be investigated for general complex c . In the case $c = -2$ there is a closed form known, $M^{[s]} = 2 \cosh(2^s \operatorname{acosh}(x/2))$. [Of course, $x^2 - 2 = 2 \cosh(2 \operatorname{acosh}(x/2))$ is essentially the double-angle formula for cosines.] And of course in the case $c = 0$ the closed form is $M^{[s]} = x^{2^s}$. For other values of c , no closed form is known, and it is likely that there is none (but that must be explained).

So, let c be a fixed real number, and write $M(x) = x^2 + c$. Use ratio set $\mu = \{\mu_0, \mu_1, \mu_2, \mu_3\}$,

$$\mu_0 = \frac{1}{\log x}, \quad \mu_1 = x^{-1}, \quad \mu_2 = e^{-x}, \quad \mu_3 = e^{-e^x}.$$

Begin with $M(x) = x^2 + c$. Then

$$M_1 := \log \circ M \circ \exp = \log(e^{2x} + c) = \log(e^{2x}(1 + ce^{-2x})) = 2x - \sum_{j=1}^{\infty} \frac{(-1)^j c^j \mu_2^{2j}}{j}.$$

The series in powers of μ_2 . Next,

$$M_2 := \log \circ M_1 \circ \exp = \log\left(2e^x - \sum_{j=1}^{\infty} \frac{(-1)^j c^j \mu_3^{2j}}{j}\right) = \log\left(2e^x \left(1 - \sum_{j=1}^{\infty} \frac{(-1)^j c^j \mu_2 \mu_3^{2j}}{2j}\right)\right).$$

Writing A for the series (in powers of μ_2, μ_3),

$$M_2 = x + \log 2 - \sum_{j=1}^{\infty} \frac{A^j}{j} = x + \log 2 + \frac{c}{2} \mu_2 \mu_3^2 - \frac{c^2}{4} \mu_2 \mu_3^4 - \frac{c^2}{8} \mu_2^2 \mu_3^4 + O(\mu_2 \mu_3^6).$$

The O term represents $\mu_2 \mu_3^6$ times a series in μ_2, μ_3 with nonnegative exponents.

Note that M_2 is *deep* in the sense of Definition 3.2, so the iterates will be computed using Abel's Equation. The solution V of the Abel equation

$$V \circ M_2 = V + \log 2$$

is found by iteration $V_0 = x$, $V_{n+1} = V_n \circ M_2 - \log 2$. The result is

$$V = x + \frac{c}{2} \mu_2 \mu_3^2 + \frac{c - c^2}{4} \mu_2 \mu_3^4 - \frac{c^2}{8} \mu_2^2 \mu_3^4 - \frac{c^2}{2} \mu_2 \mu_3^6 - \frac{c^2}{8} \mu_2^2 \mu_3^6 + O(\mu_2 \mu_3^8).$$

The O is a series in μ_2, μ_3 . For $c = -2$, closed form is:

$$V = \log \operatorname{acosh} \frac{1}{2} e^{\epsilon^x}.$$

The inverse is computed as in [10, Prop. 4.19]:

$$\begin{aligned} V^{[-1]} = x - \frac{c}{2} \mu_2 \mu_3^2 + \frac{c - c^2}{4} \mu_2 \mu_3^4 - \frac{c^2}{8} \mu_2^2 \mu_3^4 + \frac{c^3 - 3c^2}{6} \mu_2 \mu_3^6 \\ + \frac{c^3 - c^2}{8} \mu_2^2 \mu_3^6 + \frac{c^3}{24} \mu_2^3 \mu_3^6 + O(\mu_2 \mu_3^8). \end{aligned}$$

The iteration group is then a computation; for any real s ,

$$M_2^{[s]} = V^{[-1]}(V(x) + s \log 2).$$

For a fixed s , augment our ratio set with

$$\mu_4 = x^{-2^s}, \quad \mu_5 = e^{-2^s x}, \quad \mu_6 = e^{-2^s e^x}.$$

Then

$$\begin{aligned} M_2^{[s]} = x + s \log 2 + \frac{c}{2} \mu_2 \mu_3^2 - 2^{-1-s} c \mu_2 \mu_6^2 + \frac{c - c^2}{4} \mu_2 \mu_3^4 - \frac{c^2}{8} \mu_2^2 \mu_3^4 \\ + \frac{c^2}{2} \mu_2 \mu_3^2 \mu_6^2 + 2^{-2-s} c^2 \mu_2^2 \mu_3^2 \mu_6^2 + O(\mu_2 \mu_3^6 + \mu_2 \mu_6^4). \end{aligned}$$

Which of the two terms in the O is larger depends on the value of s . The relative sizes of the terms shown also depend on the value of s . If $s > 0$, then $\mu_3 \succ \mu_6$, so $M_2^{[s]} = x + s \log 2 + (c/2)\mu_2\mu_3^2 + \dots$. If $s < 0$, then $\mu_3 \prec \mu_6$, so $M_2^{[s]} = x + s \log 2 - 2^{-1-s}c\mu_2\mu_6^2 + \dots$.

Continue:

$$\begin{aligned} M_1^{[s]} = \exp \circ M_2^{[s]} \circ \log &= 2^s x + 2^{-1+s} c \mu_2^2 - \frac{c}{2} \mu_5^2 + 2^{-2+s} (c - c^2) \mu_2^4 + 2^{-1+s} c^2 \mu_2^2 \mu_5^2 \\ &+ (2^{-2+s} (c^2 - c^3) - 2^{-2+2s} c^3) \mu_2^4 \mu_5^2 - \frac{c^2 + c}{4} \mu_5^4 + O(\mu_2^6 + \mu_5^6). \end{aligned}$$

This is a series in μ_1, μ_2, μ_5 . The coefficients involve rational numbers and powers of 2^s . I do not know if μ_1 actually appears: up to this point, all terms with μ_1 cancel. Next,

$$\begin{aligned} M^{[s]} = \exp \circ M_1^{[s]} \circ \log &= x^{2^s} \left(1 + 2^{-1+s} c \mu_1^2 - \frac{c}{2} \mu_4^2 + (2^{-2+s} (c - c^2) + 2^{-3+2s} c^2) \mu_1^4 \right. \\ &\left. + 2^{-2+s} c^2 \mu_1^2 \mu_4^2 + O(\mu_1^6 + \mu_4^4) \right). \end{aligned}$$

This is a series in μ_0, μ_1, μ_4 , but I do not know if $\mu_0 = 1/\log x$ actually appears. The relative size of the terms depends on the value of s .

Let us substitute a few example values of s into this transseries:

$$\begin{aligned} M^{[1]} &= x^2 (1 + cx^{-2} + O(x^{-6})) = x^2 + c + O(x^{-4}) \\ M^{[-1]} &= x^{1/2} \left(1 - \frac{c}{2} x^{-1} + O(x^{-2}) \right) = x^{1/2} - \frac{c}{2} x^{-1/2} + O(x^{-3/2}) \\ M^{[1/2]} &= x^{\sqrt{2}} + 2^{-1/2} c x^{\sqrt{2}-2} - \frac{c}{2} x^{-\sqrt{2}} + \left(2^{-3/2} (c - c^2) + \frac{c^2}{4} \right) x^{\sqrt{2}-4} \\ &\quad + 2^{-3/2} c^2 x^{-2-\sqrt{2}} + O(x^{-3\sqrt{2}}). \end{aligned}$$

In case $c = -2$ we have:

$$M^{[1/2]} = x^{\sqrt{2}} - \sqrt{2} x^{\sqrt{2}-2} + x^{-\sqrt{2}} + \left(1 - \frac{3}{\sqrt{2}} \right) x^{\sqrt{2}-4} + \sqrt{2} x^{-2-\sqrt{2}} + O(x^{-3\sqrt{2}}),$$

which does match the transseries for the closed form

$$M^{[1/2]} = 2 \cosh \left(\sqrt{2} \operatorname{acosh} \frac{x}{2} \right).$$

Figure 1 illustrates the support of $M^{[s]}$ depending on s . The support of $M^{[s]}$ consists of certain monomials of the form x^a , where points (s, a) are shown in the figure. I have assumed that logarithmic factors are, indeed, missing. Or perhaps we could say: any monomials with logarithmic factors differ only infinitesimally from the terms shown, so even if they do exist, they make no difference in the picture.

References

- [1] M. Aschenbrenner, L. van den Dries, “Asymptotic differential algebra.” In [7], pp. 49–85




Figure 1: $M^{[s]}$ supported by monomials x^a

- [2] I. N. Baker, “Zusammensetzungen ganzer Funktionen.” *Math. Z.* **69** (1958) 121–163
- [3] A. Cayley, “On some numerical expansions.” *Quarterly Journal of Pure and Applied mathematics* **3** (1860) 366–369. Also in: *Collected Works* vol. IV, pp. 470–472
- [4] P. M. Cohn, *Universal Algebra*. Harper & Row, New York, 1965
- [5] O. Costin, “Topological construction of transseries and introduction to generalized Borel summability.” In [7], pp. 137–175
- [6] O. Costin, “Global reconstruction of analytic functions from local expansions and a new general method of converting sums into integrals.” preprint, 2007.
<http://arxiv.org/abs/math/0612121>
- [7] O. Costin, M. D. Kruskal, A. Macintyre (eds.), *Analyzable Functions and Applications* (*Contemp. Math.* **373**). Amer. Math. Soc., Providence RI, 2005
- [8] L. van den Dries, A. Macintyre, D. Marker, “Logarithmic-exponential series.” *Annals of Pure and Applied Logic* **111** (2001) 61–113

- [9] G. Edgar, “Transseries for beginners.” *Real Analysis Exchange* (to appear).
<http://arxiv.org/abs/0801.4877> or
http://www.math.ohio-state.edu/~edgar/preprints/trans_begin/
- [10] G. Edgar “Transseries: composition, recursion, and convergence.” forthcoming
<http://arxiv.org/abs/0909.1259v1> or
http://www.math.ohio-state.edu/~edgar/preprints/trans_compo/
- [11] G. Edgar, “Transseries: ratios, grids, and witnesses.” forthcoming
<http://arxiv.org/abs/0909.2430v1> or
http://www.math.ohio-state.edu/~edgar/preprints/trans_wit/
- [12] G. Edgar, “Tetration in transseries.” forthcoming
- [13] P. Erdős, E. Jabotinsky, “On analytic iteration.” *J. Analyse Math.* **8** (1960) 361–376
- [14] J. van der Hoeven, “Operators on generalized power series.” *Illinois J. Math.* **45** (2001) 1161–1190
- [15] J. van der Hoeven, *Transseries and Real Differential Algebra (Lecture Notes in Mathematics 1888)*. Springer, New York, 2006
- [16] J. van der Hoeven, “Transserial Hardy fields.” preprint, 2006
- [17] A. Korkine, “Sur un problème d’interpolation.” *Bulletin des Sciences Mathématiques et Astronomiques* (2) **6** (1882) 228–242
- [18] S. Kuhlmann, *Ordered Exponential Fields*. American Mathematical Society, Providence, RI, 2000
- [19] H. F. Wilf, *generatingfunctionology*. Academic Press, Boston, 1990.
<http://www.math.upenn.edu/~wilf/DownldGF.html>